

ORNAMENTAL LAWNS, ORNAMENTAL TURFGRASS, GOLF COURSES AND SOD FARMS - BROADCAST

General Information

GENERAL INFORMATION:

Trimec Encore Broadleaf Herbicide provides selective broadleaf control in warm season and cool season turfgrass. Established turf of warm season grasses include, but are not limited to, bermudagrass, bahiagrass, and zoysiagrass. Established turf of cool season grasses include, but are not limited to, Kentucky bluegrass, perennial ryegrass, tall fescue, and turf-type tall fescues. Do not apply this product on carpetgrass, dichondra or established turf containing desirable clovers.

USE PRECAUTIONS - OBSERVE CAREFULLY.

1. Do not pour spray solutions near these plants.
2. Do not spray on carpetgrass, dichondra nor on lawns or turf where desirable clovers are present.
3. Use only lawn type sprayers.
4. Avoid application to exposed active feeder roots of shallow-rooted trees such as poplar species and sensitive ornamentals. Do not exceed specified dosages for any area.
5. Do not use this product on or near desirable plants, including contact of spray on exposed root systems or adventitious shoots within the drip line of desirable trees and shrubs, since injury may result.
6. Do not spray bermudagrass or zoysiagrass while emerging from dormancy, unless injury can be tolerated.
7. Applications to newly seeded lawns may cause injury. Delay application until well established or after two or three mowings.
8. Do not broadcast apply when air temperatures exceed 90°F; some injury may be expected with spot treatments when air temperatures exceed 90°F.
9. Seed can be sown 3 to 4 weeks after application.
10. Do not contaminate domestic or irrigation waters.
11. Some injury can be expected to turf when used on centipedegrass or St. Augustinegrass; only spot spray weeds on these types of grasses. Clover will be damaged if sprayed.
12. After using this product, clean sprayer with soap or detergent and water, and

rinse thoroughly before applying other pesticides.

13. Failure to observe the above precautions may result in injury to desirable plants.

14. Do not apply this product through any type of irrigation system.

Limitations, Restrictions, and Exceptions

ORNAMENTAL LAWNS, ORNAMENTAL TURFGRASS, GOLF COURSES AND SOD FARMS - As Broadcast Treatments:

Apply 3.0 to 4.0 pints of product per acre per application with spray volumes of 20 to 240 gallons per acre. Maximum control of weeds will be obtained from spring or early fall applications when weeds are actively growing. Avoid applying during long excessively dry or hot periods unless adequate irrigation is available. Do not apply to newly seeded turf until well-established or after the second or third mowing. Grass seed can be sown 3 to 4 weeks after application at the recommended rates.

Limitations on broadcast treatments for turfgrass on all use sites: The maximum application rate is 4.0 pints of product per acre per application. The maximum number of broadcast applications is limited to 2 per year with a minimum of 30 days between applications. The maximum seasonal rate is 8.0 pints of product per acre per year, excluding spot treatments.

Limitations on spot treatments for all use sites: Spot treatment is defined as a treatment area no greater than 1,000 sq.ft. per acre. The maximum application rate is 1.5 fl.oz. per 1,000 sq.ft. per application. The maximum number of spot treatments is limited to 2 per year with a minimum of 30 days between applications.

Method

[Broadcast](#)

Rates

[field_rates 0](#)

-

Restricted Entry Interval

48 hours

Timings

[From spring or early fall applications when weeds are actively growing.](#)