

WOODY PLANT CONTROL - HARD-TO-CONTROL

General Information

GENERAL INFORMATION

Alligare Everett Herbicide will control many species of woody plants, annual and perennial broadleaf weeds, growing on rangeland, permanent grass pastures, CRP, fence rows, nonirrigation ditch banks, roadsides, other non-crop areas, and industrial sites.

GENERAL COMMENTS AND RESTRICTIONS

- For use on plants in non-crop and non-timber areas only. Do not apply to crops, timber, or other plants being grown for sale or other commercial use, or for commercial seed production, or for research purposes.
- Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application.
- Do not allow worker entry into areas until sprays have dried, unless applicator and other handler PPE is worn.
- The state of Arizona has not approved Alligare Everett Herbicide for use on plants grown for commercial production; specifically forests grown for commercial timber production, or on designated grazing areas.
- This product may not be applied to forage that is to be cut and sold for commercial purposes.
- Chemigation is prohibited. Do not apply through any type of irrigation system.
- Optimal control is obtained when foliar sprays are applied during warm weather when target brush and weeds are actively growing.
- Applications made under drought stress conditions will result in reduced control.

- Use low spray pressures to minimize spray drift.
- Do not use on bentgrass.
- Do not use on newly seeded grasses until grass has established a good root system and is tillering.
- Do not reseed pastures within a minimum of three weeks after treatment.
- Do not spray pastures containing desirable broadleaf forbs, especially legumes such as clover, unless injury or loss of such plants can be tolerated. However, the stand and growth of established grasses usually is improved, particularly when rainfall is adequate and grazing is deferred.
- Excessive amounts of this herbicide in the soil may temporarily inhibit seed germination and plant growth.
- Do not enter or allow others to enter the treated area until sprays have dried.

GRAZING AND HAYING RESTRICTIONS

Except for lactating dairy animals, there are no grazing restrictions following application of this product.

Grazing Lactating Dairy Animals:

- Do not allow lactating dairy animals to graze treated areas until the next growing season following application of this product.
- Do not harvest hay for 14 days after application.
- Grazed areas of non-cropland and forestry sites may be spot treated if they comprise no more than 10% of the total grazable area.

Slaughter Restrictions:

- During the season of application, withdraw livestock from grazing treated grass at least 3 days before slaughter.

Susceptible Plants

Do not apply under circumstances where spray drift may occur to food, forage, or

other plantings that might be damaged or crops thereof rendered unfit for sale, use or consumption.

Susceptible crops include, but are not limited to cotton, okra, flowers, grapes (in growing stage), fruit trees (foliage), soybeans (vegetative stage), ornamentals, sunflowers, tomatoes, beans, and other vegetables, or tobacco. Small amounts of spray drift that might not be visible may injure susceptible broadleaf plants.

Limitations, Restrictions, and Exceptions

WOODY PLANT CONTROL

Harder-To-Control Species:

High-volume applications, 1.5% mixture, conventional basal or dormant stem applications are recommended. A broadcast rate of 4 quarts per acre will increase the degree of control of these species.

Method

[Broadcast/Foliar Air](#)

[Broadcast/Foliar Ground](#)

Rates

[field rates 0](#)

-

Restricted Entry Interval

12 hours

Timings

[Postemergence \(Weed\)](#)