

# **HOUSE PLANTS - INDOORS: TO PROTECT PLANTS IN RESIDENTIAL GREENHOUSES AND FOR INDOOR LANDSCAPING**

## General Information

### DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Not for use on plants being grown for sale or for other commercial use, or for commercial seed

production or for research purposes.

Do not apply this product in enclosed areas.

Do not use food utensils to mix or apply this product.

Keep people and pets off treated areas until the spray has dried.

### SHAKE WELL BEFORE USING

#### THREE STEPS TO MIX AND APPLY:

1. For lawns and vegetables, measure the area to be sprayed. Calculate square feet by

multiplying length by width. For all other applications, apply as a thorough cover spray.

2. Use a clean sprayer. Carefully measure and mix the amount of concentrate and water as directed.

3. Spray as directed. Thorough coverage is important. Flush sprayer with clean water after

each use. Make applications when insects first appear. Repeat applications as directed.

Use Dilution rate as indicated:

1 fl.oz. = 6 tsp. or 2 Tbsp.

8 fl.oz. =  $\frac{1}{2}$  pt. or 1 cup

## Limitations, Restrictions, and Exceptions

### HOUSE PLANTS

### INDOORS: TO PROTECT PLANTS IN RESIDENTIAL GREENHOUSES AND FOR INDOOR

## LANDSCAPING:

Mix 1/4 fl.oz. in 1 gal. of water. Contact and residual: Use sufficient spray to obtain full coverage of all foliage. If application is to be made indoors, remove infested plants from living areas prior to application. Use in an area not likely to be occupied by people or pets(particularly fish). After spraying, plants can be returned to original location. Do not use more product than recommended, as this may cause some plant injury. Use with care on ferns, palms and fuchsia. As with all plants listed, especially carnations, chrysanthemums, and roses, there exists a wide variety of species and cultivars, and therefore, a wide variety of susceptibility to injury. To ascertain possible phytotoxicity, treat a few plants as described above, wait 24 hours, and check for possible injury.

### Note

Velvet plants - Fungus gnats

### Method

[Spray](#)

### Rates

[field rates 0](#)

- 

### Timings

[When insects first appear.](#)