

CONTROL WILD ONIONS AND WILD GARLIC IN LAWNS AND GRASS AREAS ALONG HIGHWAYS (NOT PASTURE OR CROPLAND)

General Information

DIRECTIONS FOR USE

APPLICATION NOTES (For All Applications):

1. Always read and follow dosage recommendations carefully and spray fields uniformly.
2. Uneven applications may result in “elephant hide” and/or bud end cracking of the Potato tubers.
3. To obtain the desired results, the plants/crop should not be stressed from either too much or too little water before or after application.
4. Apply only one application per season.
5. Do not apply to Potato plants grown for seed.
6. Rain or overhead irrigation occurring within 24 hours after application may reduce effectiveness. Spray Onions or Potatoes at least 24 hours before rain or overhead irrigation.
7. A minimum of 2 weeks should be allowed between the use of this product on Potatoes and the use of a vine desiccant.
8. This product is non-volatile, but regardless, no drift should be permitted onto nearby crops. Special care should be taken to prevent drift when aerial applications are made.
9. This product should be applied at temperatures below 85°F or at temperatures of 80°F if expected to exceed 85°F later that same day. This product is water-soluble and under high temperatures and low humidity conditions it will crystallize on the leaf surface preventing normal absorption into the plant.
10. Spray adjuvants are not recommended for application of this product on

Potatoes. For spray adjuvant use with Onions, see that specific section.

11. Although this product is compatible with most commonly used fungicides and insecticides, separate applications are recommended.

12. Occasionally, injury to the terminal growth of Potato foliage has been observed following application of this product according to label recommendations. Trials and usage have indicated that this injury does not adversely affect the yield or quality of the Potato tubers at harvest.

13. For control of wild Onions and Garlic do not apply to young seedlings less than one season old. Do not pasture livestock in, or cut hay from, treated areas.

14. This product is not corrosive to spray equipment.

15. Rinse spray equipment with water after application. Do not contaminate water by cleaning of equipment or disposal of waste.

Limitations, Restrictions, and Exceptions

TO CONTROL WILD ONIONS AND WILD GARLIC IN LAWNS AND GRASS AREAS ALONG HIGHWAYS (NOT PASTURE OR CROPLAND):

Apply 1.33 to 2.67 gallons of this product in 20 to 100 gallons of water per acre in Spring or Fall when about 90 to 95% of the bulbs have started regrowth. Use on lawns only when grass is brown and dormant. Do not use on home lawns except along edges.

TO INHIBIT GRASS GROWTH: Apply this product to green grass since brown grass will not absorb the spray. Use of this product is limited to one application per year made either in the Fall or the Spring followed by correct maintenance procedures to produce required turf growth control.

Method

[Broadcast/Foliar Air](#)

[Broadcast/Foliar Ground](#)

Rates

[field_rates 0](#)

-

Restricted Entry Interval

12 hours

Timings

In Spring or Fall when about 90 to 95% of the bulbs have started regrowth.