

## **NON-CROPLAND (EXCLUDING PUBLIC LAND) - CEREAL GRAINS - SWEET CORN (FOLIAR)**

### General Information

#### GENERAL DIRECTIONS FOR USE

Thorough crop coverage is necessary for both initial and residual control. Apply by ground in at least 10 gals./A or by air in at least 2 gals./A using sufficient water to obtain full coverage of foliage unless the label specifies otherwise. In situations where foliage is dense or pest pressure is high (heavier insect or egg pressure, larger larval stages), control can be improved by use of higher application volumes and/or higher use rates.

For cutworm control, LAMBDA-CY EC INSECTICIDE-RUP may be applied before, during, or after planting. When making soil incorporated applications, use higher rates for better control.

#### Resistance Management

LAMBDA-CY EC INSECTICIDE-RUP is a Group 3 Insecticide (contains the active ingredient lambda-cyhalothrin). Some insects are known to develop resistance to products used repeatedly for control. Because the development of resistance cannot be predicted, the use of this product should conform to resistance management strategies established for the use area. Consult your local or state agricultural authorities for details.

If resistance to this product develops in your area, this product, or other products with a similar mode of action, may not provide adequate control. If poor performance cannot be attributed to improper application or extreme weather conditions, a resistant strain of insect may be present. If you experience difficulty with control and resistance is a reasonable cause, immediately consult your local company representative or agricultural advisor for the best alternative method of control for your area.

#### Limitations, Restrictions, and Exceptions

## NON-CROPLAND (EXCLUDING PUBLIC LAND)

- See crop instructions in sections above for specific pest and rate information

### Instructions:

Spray non-cropland adjacent to agricultural areas to control insects which may migrate to and threaten crops. Follow the General Directions for Use instructions, application rates, and spray recommendations found elsewhere on the label for the adjacent crop outlet and target pests.

When foliage is dense/large, insect populations are high or larval stages are large, use the highest labeled rate for that crop-pest combination.

Repeat as necessary to maintain control.

Do not graze livestock in treated areas.

For control of adult corn rootworm beetles (*Diabrotica* spp.) as part of an aerial applied corn rootworm control program use a minimum of 0.025 lb. a.i. (3.2 fl. oz.)/A.

- Aphid spp: Suppression only. See resistance statement under General Directions for Use.

- Armyworm, Fall Armyworm, Southern Armyworm, Yellowstriped Armyworm: Use higher rates for large larvae.

- Spider Mite spp: Suppression only.

- Beet Armyworm: Use higher rates for large larvae. See resistance statement under General Directions for Use.

### Method

[Broadcast/Foliar Air](#)

[Broadcast/Foliar Ground](#)

### Rates

[field\\_rates 0](#)

[field\\_rates 1](#)

•

### Restricted Entry Interval

24 hours

Timings

[When insect populations are high or larval stages are large.](#)