

# **CRANBERRY**

## General Information

### DIRECTIONS FOR USE

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

MANZATE PRO-STICK™, a dispersible granule containing mancozeb, is labeled for use as a spray for the control of many important plant diseases.

### Application Instructions

AS A SPRAY (Ground or Aerial Equipment) - Apply MANZATE PRO-STICK at the rate shown; use sufficient water to provide thorough coverage: use 20 to 100 gallons per acre for ground equipment and no less than 2 gallons per acre for aircraft. Add MANZATE PRO-STICK slowly to water in the spray tank with agitation, or premix thoroughly in separate holding tank for concentrate or aircraft sprayers. Continuous agitation is required to keep the product in suspension. A spreader-sticker spray adjuvant may be used with this product if needed; contact your local product distributor or United Phosphorus, Inc. representative for specific recommendations. If tank mixed, follow more restrictive labeling of any tank mix partner. Do not tank mix with any product that contains a prohibition on tank mixing.

### Restrictions

#### Foliar Applications

Where EBDC Products Used Allow the Same Maximum Poundage of Active Ingredient Per Acre Per Season

If more than one product containing an EBDC active ingredient (maneb, mancozeb or metiram) is used on a crop during the same growing season and the EBDC products used allow the same maximum poundage of active ingredient per acre per season, then the total poundage of all such EBDC products used must not exceed

any one of the specified individual EBDC product maximum seasonal poundage of active ingredient allowed per acre.

#### Where EBDC Products Used Allow Different Maximum Poundage of Active Ingredient Per Acre Per Season

If more than one product containing an EBDC active ingredient is used on a crop during the same growing season and the EBDC products used allow different maximum poundage of active ingredient per acre per season, then the total poundage of all such EBDC products used must not exceed the lowest specified individual EBDC product maximum seasonal poundage of active ingredient allowed per acre.

#### Seed Treatment

In addition to the maximum number of foliar applications permitted by the formula stated above, a single application for seed treatment may be made on crops which have registered seed treatment uses.

#### Chemigation

Apply MANZATE PRO-STICK fungicide only through sprinkler systems including center pivot, lateral move, end tow, side (wheel) roll, traveler, big gun, solid set or hand move irrigation systems. Do not apply MANZATE PRO-STICK through any other type of irrigation system.

Crop injury, lack of effectiveness, or illegal pesticide residues in the crop can result from non-uniform distribution of treated water.

If you have questions about calibration, you should contact State Extension Service Specialists, equipment manufacturers or other experts.

Do not connect an irrigation system (including greenhouse systems) used for pesticide application to a public water system unless the pesticide label-prescribed safety devices for public water systems are in place.

A person knowledgeable of the chemigation system and responsible for its operation, or under the supervision of the responsible person, shall shut the system down and make necessary adjustments should the need arise.

## Specific Instructions for Public Water Systems:

1. Public water system means a system for the provision to the public of piped water for human consumption if such system has at least 15 service connections or regularly serves an average of at least 25 individuals daily at least 60 days out of the year.
2. Chemigation systems connected to public water systems must contain a functional, reduced-pressure zone, backflow preventer (RPZ) or the functional equivalent in the water supply line upstream from the point of pesticide introduction. As an option to the RPZ, the water from the public water system should be discharged into a reservoir tank prior to pesticide introduction. There shall be a complete physical break (air gap) between the outlet end of the fill pipe and the top or overflow rim of the reservoir tank of at least twice the inside diameter of the fill pipe.
3. The pesticide injection pipeline must contain a functional, automatic, quick-closing check valve to prevent the flow of fluid back toward the injection pump.
4. The pesticide injection pipeline must contain a functional, normally closed, solenoid-operated valve located on the intake side of the injection pump and connected to the system interlock to prevent fluid from being withdrawn from the supply tank when the irrigation system is either automatically or manually shut down.
5. The system must contain functional interlocking controls to automatically shut off the pesticide injection pump when the water pump motor stops, or in cases where there is no water pump, when the water pressure decreases to the point where pesticide distribution is adversely affected.
6. Systems must use a metering pump, such as a positive displacement injection pump (e.g., diaphragm pump) effectively designed and constructed of materials that are compatible with pesticides and capable of being fitted with a system interlock.
7. Do not apply when wind speed favors drift beyond the area intended for treatment.

## Specific Instructions for Sprinkler Irrigation Systems:

1. The system must contain a functional check valve, vacuum relief valve, and low pressure drain appropriately located on the irrigation pipeline to prevent water source contamination from backflow.

2. The pesticide injection pipeline must contain a functional, automatic, quick-closing check valve to prevent the flow of fluid back toward the injection pump.
3. The pesticide injection pipeline must also contain a functional, normally closed, solenoid-operated valve located on the intake side of the injection pump and connected to the system interlock to prevent fluid from being withdrawn from the supply tank when the irrigation system is either automatically or manually shut down.
4. The system must contain functional interlocking controls to automatically shut off the pesticide injection pump when the water pump motor stops.
5. The irrigation line or water pump must include a functional pressure switch which will stop the water pump motor when the water pressure decreases to the point where pesticide distribution is adversely affected.
6. Systems must use a metering pump, such as a positive displacement injection pump (e.g., diaphragm pump) effectively designed and constructed of materials that are compatible with pesticides and capable of being fitted with a system interlock.
7. Do not apply when wind speed favors drift beyond the area intended for treatment.
8. Good agitation is required in the injection tank.
9. In moving systems, apply specified dosage of MANZATE PRO-STICK as a continuous injection. In non-moving systems inject MANZATE PRO-STICK for 15 to 30 minutes at end of cycle. Use the least amount of water possible consistent with uniform coverage.
10. Mix the amount of MANZATE PRO-STICK needed for acreage to be treated into the quantity of water determined during prior calibration. For moving systems inject into the system continuously for one complete revolution of the field. For non-moving systems inject into system for the time established during calibration.
11. Stop injection equipment after treatment is completed and continue to operate irrigation equipment until all MANZATE PRO-STICK is flushed from system.

### Spray Drift Management

A variety of factors including weather conditions (e.g. wind direction, wind speed, temperature, relative humidity) and method of application (e.g. ground, aerial, airblast, chemigation) can influence pesticide drift. The applicator must evaluate all factors and make appropriate adjustments when applying this product.

### Wind Speed

Do not apply at wind speeds greater than 15 mph.

### Temperature Inversions

If applying at wind speeds less than 3 mph, the applicator must determine if a) conditions of temperature inversion exist, or b) stable atmospheric conditions exist at or below nozzle height. Do not make applications into areas of temperature inversions or stable atmospheric conditions.

### Other State and Local Requirements

Applicators must follow all state and local pesticide drift requirements regarding application of mancozeb. Where states have more stringent regulations, they must be observed.

### Equipment

All aerial and ground application equipment must be properly maintained and calibrated using appropriate carriers or surrogates.

### Additional requirements for aerial applications:

1. The boom length must not exceed 75% of the wingspan or 90% of the rotor blade diameter.
2. Release spray at the lowest height consistent with efficacy and flight safety. Do not release spray at a height greater than 10 feet above the crop canopy unless a greater height is required for aircraft safety.
3. When applications are made with a crosswind, the swath must be displaced downwind. The applicator must compensate for this displacement at the up and downwind edge of the application area by adjusting the path of the aircraft upwind.

### Additional requirements for ground boom application:

1. Do not apply with a nozzle height greater than 4 feet above the crop canopy.

### Limitations, Restrictions, and Exceptions

#### CRANBERRY

#### Timing/Intervals

- Also refer to Directions for Use
- Start applications at mid-bloom and repeat at 7 to 10 day intervals.

## Restrictions/Comments

- Do not apply more than 18 lbs (13.5 lbs active) per acre per season.

## Method

[Broadcast/Foliar Air](#)

[Broadcast/Foliar Ground](#)

## Pre-Harvest Interval

30 days

## Rates

[field\\_rates 0](#)

- 

## Restricted Entry Interval

24 hours

## Timings

[At mid-bloom.](#)