

SUPPRESSION OF PEAR PSYLLA NYMPHS ON PEARS - OREGON

General Information

Application Instructions

AS A SPRAY (Ground or Aerial Equipment) - Apply MANZATE PRO-STICK at the rate shown; use sufficient water to provide thorough coverage: use 20 to 100 gallons per acre for ground equipment and no less than 2 gallons per acre for aircraft. Add MANZATE PRO-STICK slowly to water in the spray tank with agitation, or premix thoroughly in separate holding tank for concentrate or aircraft sprayers. Continuous agitation is required to keep the product in suspension. A spreader-sticker spray adjuvant may be used with this product if needed; contact your local product distributor or United Phosphorus, Inc. representative for specific recommendations. If tank mixed, follow more restrictive labeling of any tank mix partner. Do not tank mix with any product that contains a prohibition on tank mixing.

Restrictions

Foliar Applications

Where EBDC Products Used Allow the Same Maximum Poundage of Active Ingredient Per Acre Per Season

If more than one product containing an EBDC active ingredient (maneb, mancozeb or metiram) is used on a crop during the same growing season and the EBDC products used allow the same maximum poundage of active ingredient per acre per season, then the total poundage of all such EBDC products used must not exceed any one of the specified individual EBDC product maximum seasonal poundage of active ingredient allowed per acre.

Where EBDC Products Used Allow Different Maximum Poundage of Active Ingredient Per Acre Per Season

If more than one product containing an EBDC active ingredient is used on a crop during the same growing season and the EBDC products used allow different maximum poundage of active ingredient per acre per season, then the total poundage of all such EBDC products used must not exceed the lowest specified

individual EBDC product maximum seasonal poundage of active ingredient allowed per acre.

Seed Treatment

In addition to the maximum number of foliar applications permitted by the formula stated above, a single application for seed treatment may be made on crops which have registered seed treatment uses.

Limitations, Restrictions, and Exceptions

APPLICATION INSTRUCTIONS

Apply MANZATE PRO-STICK at the rate of 6 pounds per acre as a thorough coverage spray.

Begin applications at early bloom (pink bud stage) and discontinue after petal fall.

Maximum of 2 applications for this use and a minimum interval between applications of 7 days.

RESTRICTIONS

- Do not combine or integrate this use with the "Extended Application Schedule" found on the federally approved label for MANZATE PRO-STICK.
- Do not apply after petal fall.
- Do not graze livestock in treated areas.

Method

[Spray](#)

Rates

[field rates 0](#)

•

Restricted Entry Interval

24 hours

Timings

[At early bloom \(pink bud stage\) and discontinue after petal fall.](#)