

# **USE ON PEARS FOR SUPPRESSION OF PEAR PSYLLA NYMPHS - WASHINGTON**

## General Information

## Application Instructions

AS A SPRAY (Ground or Aerial Equipment) - Apply MANZATE PRO-STICK at the rate shown; use sufficient water to provide thorough coverage: use 20 to 100 gallons per acre for ground equipment and no less than 2 gallons per acre for aircraft. Add MANZATE PRO-STICK slowly to water in the spray tank with agitation, or premix thoroughly in separate holding tank for concentrate or aircraft sprayers. Continuous agitation is required to keep the product in suspension. A spreader-sticker spray adjuvant may be used with this product if needed; contact your local product distributor or United Phosphorus, Inc. representative for specific recommendations. If tank mixed, follow more restrictive labeling of any tank mix partner. Do not tank mix with any product that contains a prohibition on tank mixing.

## Restrictions

### Foliar Applications

#### Where EBDC Products Used Allow the Same Maximum Poundage of Active Ingredient Per Acre Per Season

If more than one product containing an EBDC active ingredient (maneb, mancozeb or metiram) is used on a crop during the same growing season and the EBDC products used allow the same maximum poundage of active ingredient per acre per season, then the total poundage of all such EBDC products used must not exceed any one of the specified individual EBDC product maximum seasonal poundage of active ingredient allowed per acre.

#### Where EBDC Products Used Allow Different Maximum Poundage of Active Ingredient Per Acre Per Season

If more than one product containing an EBDC active ingredient is used on a crop during the same growing season and the EBDC products used allow different maximum poundage of active ingredient per acre per season, then the total

poundage of all such EBDC products used must not exceed the lowest specified individual EBDC product maximum seasonal poundage of active ingredient allowed per acre.

## Seed Treatment

In addition to the maximum number of foliar applications permitted by the formula stated above, a single application for seed treatment may be made on crops which have registered seed treatment uses.

## Limitations, Restrictions, and Exceptions

### FOR USE ON PEARS FOR SUPPRESSION OF PEAR PSYLLA NYMPHS

Application rate and timing: Apply 8 pounds of product (6 lb ai) per acre. Begin thorough coverage applications at early bloom (pink) or at first signs of psylla activity. Reapply as needed on a 7- to 10-day interval and discontinue after petal fall. Do not apply more than three applications per acre per season.

Method of application: Apply as a broadcast foliar spray by ground equipment calibrated to deliver a minimum of 50 gallons per acre to trees 10 feet tall or less. For trees greater than 10 feet use a minimum of 100 gallons per acre. See package label for use directions and restrictions for ground applications.

### Restrictions / Precautions:

- Do not combine or integrate this use with the "Extended Application Schedule" found on the federally approved label.
- Do not apply after petal fall.
- Do not graze livestock in treated areas.
- For use under this SLN label, do not apply this product using aerial application equipment or through any type of irrigation system.

## Method

### [Broadcast/Foliar Ground](#)

### Pre-Harvest Interval

77 days

Rates

[field\\_rates 0](#)

[field\\_rates 1](#)

- 

Restricted Entry Interval

24 hours

Timings

[At early bloom \(pink\) or at first signs of psylla activity.](#)