

# FOLIAR APPLICATIONS - NUTS AND POME FRUIT

## General Information

### General Information

BRANDT LIQUID ZINC 10% is recommended to prevent and correct zinc deficiencies of field, row, vegetable, fruit, tree, and vine crops. Applied alone, with pesticides, or with other fertilizers, this product may be applied directly to the soil or to the foliage of growing crops via ground sprayers, aircraft or fertigation systems.

### Mixing and Handling Instructions

Apply in sufficient water for thorough coverage. For aerial applications, use at least 1 gallon of water per pint of product.

Put 1/3 to 2/3 of total required water volume in tank. Add recommended amount of this product before adding pesticides, adjuvants, or other fertilizers. Agitate until thoroughly mixed, then fill tank with remainder of water.

BRANDT LIQUID ZINC 10% is compatible with most pesticides and fertilizers. However, if the desired combination has not been previously used, check for chemical mixture compatibility using a jar test prior to application. When mixing with high polyphosphate fertilizers, do not exceed a phosphate:zinc ratio of 10:1. Do not tank mix with carbamate pesticides as efficacy will decrease.

### Limitations, Restrictions, and Exceptions

FOLIAR APPLICATION: Apply in the early morning or late evening for optimum results. Over application may be phytotoxic or cause undesirable foliage effects.

Nuts and Pome Fruit: Apply at green tip, and thereafter at ~30 day intervals, or as needed.

### Method

[Broadcast/Foliar Air](#)

[Broadcast/Foliar Ground](#)

### Rates

[field rates 0](#)

-

Timings

N. A.