

FRUIT ELIMINATION (ORNAMENTAL USE ONLY) - CAROB

General Information

INFORMATION

This product: eliminates leafy mistletoe from ornamental deciduous trees and dwarf mistletoe from ornamental conifers. It also removes unwanted fruit on ornamental apple, crabapple, carob, elm, flowering pear, flowering plum, horsechestnut (Buckeye), oak, olive, sour orange, sweetgum, and sycamore trees.

Contact your Extension Pomologist, Farm Advisor, Horticultural Specialist or United Phosphorus Representative for local requirements on product spray volume, spray equipment and rates of application for varying weather conditions.

APPLICATION VOLUMES AND SPRAY COVERAGE

For optimum product efficacy, thorough spray coverage is necessary. This can be influenced by type of spray equipment, spray boom setup, nozzle selection, plant size, canopy density and spray pressure. Depending on these choices, the necessary spray volume will vary. For applications by air in California and Arizona, more than 5 gallons of water per acre should be used.

USE PRECAUTIONS

DO NOT MIX THIS PRODUCT WITH AMMONIUM THIOSULFATE AS IT MAY CREATE TOXIC FUMES. Other than permitted on the label, this product should not be used with additives.

Upon mixture, this product should be applied as soon as possible; in no case should the spray solution be stored overnight. Detrimental changes to plant growth, reduced yields, and plant injury may result from spray drift of this product to nearby crops and thus should be avoided. Do not plant another crop within 30 days after treatment.

This product is corrosive. Therefore, spills of concentrated product on the aircraft or other spray equipment should be avoided. Should such contact be made, immediately rinse with water.

Limitations, Restrictions, and Exceptions

FRUIT ELIMINATION

(Ornamental Use Only)

UNDESIRABLE FRUIT ELIMINATION: A foliar spray of Phoenix Cardinal will reduce or eliminate undesirable fruit development on apple, crabapple, carob, elm, flowering pear, flowering plum, horsechestnut (Buckeye), oak, olive, sour orange, sweetgum, and sycamore trees.

Instructions

Apply as a foliar spray at the mid to full bloom stage, prior to fruit set.

Wet foliage thoroughly.

Over application of Phoenix Cardinal can result in excessive leaf drop and/or tree defoliation.

Comments

Application must be made before fruit set for best results.

Do not treat weak trees or trees under stress (drought, insect or disease damaged trees) as excessive leaf drop or twig drop can result.

Some leaf drop or temporary leaf yellowing may occur after treatment.

Do not use on small red fruited varieties of crabapple as fruit elimination will not be satisfactory.

Method

[Foliar spray](#)

Rates

[field rates 0](#)

-

Restricted Entry Interval

48 hours

The REI is 72 hours in areas where average rainfall is less than 25 inches per year.

Timings

[At the mid to full bloom stage, prior to fruit set.](#)