

CHERRIES - SWEET - SINGLE APPLICATION

General Information

GENERAL INFORMATION

- Use only as directed. Read and thoroughly understand the label before making applications.
- The term 'grams of active ingredient per acre' is represented by 'g ai/acre' throughout this product label. See conversion table at the end of Directions for Use.
- Thoroughly spray all parts of the plant or crop to obtain the desired result.
- Prepare solution concentrations by mixing the required amount of product with water only in a clean empty spray tank.
- Dispose of any unused spray material at the end of the day. Refer to the Storage and Disposal section of the label for pesticide disposal instruction.
- When a range of rates is indicated, consult your local experimental station, distributor, or agricultural extension agent for the best program suited to your local conditions.
- Falgro 20SP works best when using water of neutral or slightly acidic pH. Ensure water pH is less than 8.5.
- Absorption of Falgro 20SP into the plant is greatest under slow drying conditions. Nighttime applications will be more effective when daytime conditions cause rapid drying. Re-apply Falgro 20SP if significant rain occurs within 2 hours of application.
- Falgro 20SP has a 0-day pre harvest interval (PHI).
- Do not apply using ULV application methods. For aerial applications spray volumes must be greater than 2 gallons per acre (10 gallons per acre for tree crops).
- Consult your local experimental station, distributor, or agricultural extension agent regarding the compatibility of gibberellic acid with other compounds.

Limitations, Restrictions, and Exceptions

CHERRIES - SWEET

Application of Falgro 20SP produces larger, brighter colored and firmer fruit. This treatment may delay color development and harvest.

SINGLE APPLICATION

Make one application of 16 to 48 g ai/acre (do not exceed a total of 48 g ai/acre) as a dilute spray on large mature trees when the fruit is light green to straw colored in sufficient water to ensure thorough wetting.

Method

[Broadcast/Foliar Air](#)

[Broadcast/Foliar Ground](#)

Rates

[field_rates 0](#)

•

Restricted Entry Interval

4 hours

Timings

[When the fruit is light green to straw colored.](#)