

VEGETATION MANAGEMENT (NON-CROP AREAS)

General Information

GENERAL INFORMATION

WHERE TO USE

Alligare Prodiamine 65 WG Herbicide is a preemergence herbicide that provides residual control of many grass and broadleaf weeds in:

- Established turfgrass (excluding golf course putting greens), lawns, and sod nurseries
- Container, field-grown, and landscape ornamentals
- Conifer and hardwood seedling nurseries
- Established perennial and wildflower plantings
- Non-crop areas including managed rights-of-way for transportation systems and utilities (including roadways, roadsides, railways, and equipment yards)
- Facilities including substations, tank farms, pumping stations, parking and storage areas, and ungrazed fence rows
- Christmas tree farms
- Drainage ditches in California and Arizona only

HOW ALLIGARE PRODIAMINE 65 WG HERBICIDE WORKS

Alligare Prodiamine 65 WG Herbicide controls susceptible weeds by preventing growth and development of newly germinated weed seeds. Weed control is most effective when Alligare Prodiamine 65 WG Herbicide is activated by at least 0.5 inch of rainfall or irrigation or shallow incorporation (1 to 2 inches) before weed seeds germinate and within 14 days following application.

USE PRECAUTIONS

- Do not graze or feed livestock forage cut from areas treated with Alligare Prodiamine 65 WG Herbicide.
- Do not apply Alligare Prodiamine 65 WG Herbicide to plants that will be consumed for food use.
- Follow all applicable directions, restrictions, and precautions on the labels of EPA-registered tank mix partners.
- Do not blend Alligare Prodiamine 65 WG Herbicide onto dry fertilizer or any other granular material.
- Chemigation Statement: Do not apply this product through any type of irrigation system unless instructed otherwise in the label.
- Do not apply aerially.
- Do not apply to golf course putting greens.

NEW PLANTINGS, REPLANTING, AND ROTATIONAL PLANTINGS

Nursery, landscape, or non-crop land areas treated with Alligare Prodiamine 65 WG Herbicide should be rotated only to ornamental species listed on the label for 1 year following application unless the following test has shown species safety:

Before planting a species not listed on the label, it is recommended that several test strips of an indicator plant such as wheat, sorghum, or corn be sown into the treated area. If the indicator plants germinate and grow normally to a height of 12 inches with normal root development, it is safe to plant. In areas disturbed by new plantings or replanting of labeled species, it may be necessary to retreat exposed soil to maintain satisfactory weed control.

When an adjuvant is to be used with this product, Alligare, LLC suggests the use of a Chemical Producers and Distributors Association certified adjuvant.

APPLICATION

Apply Alligare Prodiamine 65 WG Herbicide in a minimum of 20 gals./acre (0.5 gal./1,000 sq. ft.) of carrier (water and/or fluid fertilizer) using a calibrated, low-pressure sprayer with 50 mesh or coarser screens. A broadcast boom or handheld

wand designed for herbicide or insecticide application will provide the best results. Select nozzle pressure and gallonage to provide complete coverage.

Refer in the label for tank mix information.

Limitations, Restrictions, and Exceptions

VEGETATION MANAGEMENT (NON-CROP AREAS)

- May be applied in soil surfaces for preemergence control of many grass and broadleaf weeds.

- Is most effective when activated by at least 0.5 inch rainfall or irrigation, or shallow incorporation before weed seeds germinate and within 14 days after application.

COMMENTS/ INSTRUCTIONS

Use higher rate for longer control.

This product may be applied more than once per year but do not apply more than 2.3 lbs./A per year.

Alligare Prodiamine 65 WG Herbicide may be tank mixed with other registered herbicides to provide a broader spectrum of weed control or to control emerged weeds or brush. Tank mixes with Alligare Prodiamine 65 WG Herbicide are for use only in states where the tank mix partner(s) are registered for the application site.

Method

[Surface](#)

[Spray](#)

Rates

[field rates 0](#)

[field rates 1](#)

-

Restricted Entry Interval

12 hours

Exception: If this product is soil-injected or soil-incorporated, the Worker Protection Standard, under certain circumstances, allows workers to enter the treated area if

there will be no contact with anything that has been treated.

Timings

[Preemergence \(Weed\)](#)