

CHERRY (SWEET AND TART) - POWDERY MILDEW ETC.

General Information

PRODUCT INFORMATION

LUNA SENSATION is a broad-spectrum fungicide with preventative, systemic, and curative properties labeled for the control or suppression of certain crop diseases.

LABELED USES

Almond, apple, cherry, pecan, pistachio, tree nuts, and watermelon.

APPLICATION INFORMATION

Use sufficient water volume to provide thorough and uniform coverage to obtain the most effective disease control. Do not make applications when conditions favor drift. Avoid spraying when windy, high temperature, drought, dusty, low relative humidity, or temperature inversion conditions exist.

Ground Application

For ground application equipment, a minimum of 50 gallons of water per acre for tree crops and 10 gallons of water per acre for field and vegetable crops is required.

Aerial Application

For aerial application equipment, a minimum of 15 gallons of water per acre for tree crops and 5 gallons of water per acre for field and vegetable crops is required. Do not apply by aerial application in New York State.

Air-Blast Application

Air-assisted or air-blast sprayers move spray droplets into the crop canopy using a forced-air system. The fan should be set up to deliver only enough air volume to penetrate the canopy and provide good coverage. Adjust deflectors or other aiming devices to direct spray only to the target area. Equip sprayers with nozzles that provide accurate and uniform application.

PRODUCT RESTRICTIONS AND LIMITATIONS

Do not apply more than the maximum seasonal rate for each specific crop from any combination of products containing FLUOPYRAM or TRIFLOXYSTROBIN.

Limitations, Restrictions, and Exceptions

CHERRY (SWEET AND TART)

Powdery mildew/Rusty spot: Follow leaf spot schedule until terminal growth ceases.

Cherry leaf spot: Begin application at petal fall or when first leaves unfold and continue applications at 7- to 14-day intervals. Applications should be made at 7-day intervals early in the growing season when terminal growth is rapid. A post-harvest application may be made to maintain control and reduce overwintering inoculum.

Restrictions: To limit the potential for development of disease resistance to these fungicide classes, do not make more than 2 sequential applications of LUNA SENSATION or any Group 7 or Group 11 containing fungicide before rotating with a fungicide from a different Group.

Pre-Harvest Interval

1 day

Rates

[field rates 0](#)

-