

# **INDOOR TREATMENTS - ANTS - SURFACE APPLICATION**

## General Information

### USE RESTRICTIONS

- DO NOT apply to humans, their clothing, or bedding. DO NOT allow children or pets to contact treated surfaces until spray has dried.
- DO NOT use on household pets. Cover fish tanks prior to application.
- DO NOT apply this product in patient rooms while occupied or in any rooms while occupied by the elderly or infirm.
- DO NOT apply to classrooms when in use.
- DO NOT contaminate food, feedstuffs, or water supply. DO NOT contaminate food preparation surfaces, kitchen utensils, dishes, or feed storage containers. Cover all food/feed contact surfaces and cooking utensils during treatment or thoroughly wash before using.
- DO NOT apply water based sprays of this product to conduits, motor housing, junction and switch boxes, or other electrical or electronic equipment because of possible shock hazard or damage. DO NOT spray where electrical short circuits might result, such as wall outlets, conduits, etc.
- DO NOT make outdoor applications during rain.
- DO NOT apply this product through any type of irrigation system.
- DO NOT water the treated area to the point of runoff.
- DO NOT apply directly to impervious horizontal surfaces such as sidewalks, driveways, and patios except as a spot or Crack & Crevice treatment. During application, DO NOT allow pesticide to enter or runoff into storm drains, drainage ditches, gutters, or surface waters.
- All outdoor applications, if permitted elsewhere on the label, must be limited to spot or Crack & Crevice treatments only, except for the following permitted uses, if allowed elsewhere on the label:
  1. Applications to soil or vegetation, as listed on the label, around structures;
  2. Applications to lawns, turf, and other vegetation, as listed on the label;
  3. Applications to the side of a building, up to a maximum height of 3 feet above grade;
  4. Applications to the underside of eaves, soffits, doors, or windows permanently

protected from rainfall by a covering, overhang, awning, or other structure; 5.

Applications around potential pest entry points into buildings, when limited to a surface band not to exceed 1 inch in width;

6. Applications made through the use of a coarse, low pressure spray to only those portions of surfaces that are directly above bare soil, lawn, turf, mulch, or other vegetation, as listed on the label, and not over an impervious surface, drainage, or other condition that could result in runoff into storm drains, drainage ditches, gutters, or surface waters, in order to control occasional invaders or aggregating pests.

- DO NOT apply directly to sewers or drains, or to any area like a gutter where drainage to sewers, storm drains, water bodies or aquatic habitat can occur, except as directed by the label.

## PRODUCT INFORMATION

This product is a liquid capsule suspension containing 6% cyfluthrin which will provide extended residual control of the pests listed on the label. This product may be applied using compressed air, backpack, or power spray equipment. Thoroughly clean spray equipment before using this product. Shake concentrate well before mixing. First add approximately 1/2 the required water to the tank, then add the proper amount of concentrate. Agitate, add the remaining water, and thoroughly agitate the solution again. When emptying the bottle, triple rinse with water and add the rinsate to the spray tank. Continue to agitate solution occasionally during use to ensure even coverage. Shake or reagitate solution if spraying is interrupted or if dilution is left in sprayer overnight. If spray screens are used, they should be  $\geq$  50 mesh. Water may be used at temperatures up to 160°F.

This product may be sprayed on any surface which will not be damaged or stained by water. Heavy applications may leave a visible deposit. On some surfaces this deposit can be removed with a damp cloth or sponge.

Limitations, Restrictions, and Exceptions

## SURFACE APPLICATIONS

Use 1.0 to 2.0 fluid ounces of concentrate per 1,000 square feet in sufficient water to adequately cover the area being treated without dripping or runoff. The higher rate should be used for clean-outs, for longer residual control, or for heavy insect

infestations.

Apply to walls; floors; ceilings; in and around cupboards; between, behind, and beneath equipment and appliances; around floor drains, window and door frames, and light fixtures; and on the underside of shelves, drawers, and in similar areas. Applications may also be made to floor surfaces, along walls, and around air ducts.

As a perimeter spray to help prevent pest infestations, treat a band of soil 6 to 10 feet wide around and adjacent to the building or structure. Also treat the building foundation and sides, up to a maximum height of 3 feet, where pests are active and may find entrance. Treat other insect entry points such as doors, windows, and under siding and eaves of structure.

## INDOOR TREATMENTS

For indoor pests and occasional invaders, make applications in a manner as previously described under Surface Applications and/or Spot, Crack & Crevice, or Void Applications. Pay particular attention to entry points and harborage areas.

DO NOT use in food/feed areas of food/feed handling establishments, restaurants, or other areas where food/feed is commercially prepared or processed. DO NOT use in serving areas while food/feed is exposed or facility is in operation. Serving areas include areas where prepared foods are served, such as dining rooms, and exclude areas where foods may be prepared or held. In the home, cover all food/feed processing surfaces and utensils during treatment or thoroughly wash before using. Cover or remove exposed food/feed. Non-food/feed areas are areas such as garbage rooms, rest rooms, floor drains (to sewers), entries and vestibules, offices, locker rooms, machine rooms, boiler rooms, garages, mop closets, and storage (after canning or bottling).

## ANTS

Inject cracks, crevices, and voids where ants may be traveling and/or harboring such as around doors, window frames, and other structural cracks, crevices, and voids. Spot treatment of trails and feeding areas may also be made.

Method

[Surface](#)

[Injection](#)

[Coarse Spray](#)

Rates

[field\\_rates 0](#)

- 

Timings

[N.A.](#)