

GRAPE (NOT FOR USE IN MISSISSIPPI) - SOIL SURFACE APPLICATION

General Information

Grape

(Not for use in Mississippi)

Soil Surface Application

(For use in areas east of the Continental Divide only)

Apply Lorsban Advanced just before the pest emerges from the soil.

Apply 2 quarts of the diluted spray mixture to the soil surface on a 15-square foot area (4.4 foot circle) around the base of each vine.

Specific Use Restrictions for Soil Surface Application:

- Do not allow spray to contact fruit or foliage.

Specific Use Restrictions for Soil Surface Application and Prebloom Application:

- Based upon available residue data, the use of Lorsban Advanced in grapes is restricted to areas east of the Continental Divide only. Do not use in the state of Mississippi.

Limitations, Restrictions, and Exceptions

Grape

(Not for use in Mississippi)

Soil Surface Application

(For use in areas east of the Continental Divide only)

Apply Lorsban Advanced just before the pest emerges from the soil.

Apply 2 quarts of the diluted spray mixture to the soil surface on a 15-square foot area (4.4 foot circle) around the base of each vine.

Specific Use Restrictions for Soil Surface Application:

- Do not allow spray to contact fruit or foliage.

Method

[Soil Surface](#)

Pre-Harvest Interval

35 days

Rates

[field rates 0](#)

-

Restricted Entry Interval

24 hours