

CORN (FIELD, SWEET, SEED) - POSTEMERGENCE APPLICATION - BILLBUGS, ETC.

General Information

Product Information

Lorsban Advanced insecticide is an emulsion in water for use in listed crops. This product resists washoff once it is dry. Target pests and application rates are provided in the accompanying tables.

Use Precautions and Restrictions

Insect control may be reduced at low spray volumes under high temperature and wind conditions.

Some reduction in insect control may occur under unusually cool conditions.

Flood irrigation: To avoid contamination of irrigation tail waters, do not flood irrigate within 24 hours following a soil surface or foliar application of Lorsban Advanced.

Do not aerially apply this product in Mississippi.

Insecticide Resistance Management (IRM)

Lorsban Advanced contains a Group 1B insecticide. Insect/mite biotypes with acquired resistance to Group 1B may eventually dominate the insect/mite population if Group 1B insecticides are used repeatedly in the same field or in successive years as the primary method of control for targeted species. This may result in partial or total loss of control of those species by Lorsban Advanced or other Group 1B insecticides.

To delay development of insecticide resistance:

- Avoid consecutive use of insecticides with the same mode of action (same insecticide group) on the same insect species.
- Use tank mixtures or premix products containing insecticides with different modes of action (different insecticide groups) provided the products are registered for the intended use.
- Base insecticide use upon comprehensive Integrated Pest Management (IPM)

programs.

- Monitor treated insect populations in the field for loss of effectiveness.
- Contact your local extension specialist, certified crop advisor, and or manufacturer for insecticide resistance management and/or IPM recommendations for the specific site and resistant pest problems.
- For further information or to report suspected resistance, you may contact Dow AgroSciences by calling 800-258-3033.

Application Directions

Broadcast Foliar Application

Apply with conventional power-operated spray equipment using nozzles and spray pressures specified for insecticides. Apply Lorsban Advanced in a spray volume of not less than 2 gallons per acre (gpa) for aerial application equipment (fixed wing or helicopter) or not less than 10 gpa for ground equipment, unless otherwise specified. Increase spray volume to ensure adequate coverage with increased density and height of crop canopy.

Ground Application: Orient the boom and nozzles so that uniform coverage is obtained. The swath width should not be wider than the boom. Follow nozzle manufacturer's specifications for insecticide nozzles with respect to nozzle type, pressure, and spacing.

Broadcast Soil Application

Apply with conventional power-operated spray equipment that will apply the product uniformly to the soil surface. Use nozzles that produce medium or coarse droplets (255 to 400 microns). Unless otherwise indicated, a spray volume of 10 gpa or more is needed. For band application, use proportionally less spray volume.

Aerial Application

Use a minimum spray volume of 2 gpa. Mark swaths by mechanical flagging, permanent markers or GPS equipment.

Chemigation Application

Apply Lorsban Advanced through properly equipped chemigation systems for insect control in alfalfa, almond (orchard floors only), citrus (orchard floors only), corn (field and sweet), cotton, cranberry, peppermint, sorghum, soybeans, spearmint,

sugarbeet, orchard floors (pecan and walnut), and wheat, or other crops as specified in Dow AgroSciences supplemental labeling. Do not apply this product by chemigation unless specified in crop-specific directions in this label or Dow AgroSciences supplemental labeling. Do not apply to labeled crops through any other type of irrigation system.

Note: Unless otherwise indicated in specific use directions, the application rates for chemigation are the same as those specified for broadcast application.

Limitations, Restrictions, and Exceptions

CORN (FIELD, SWEET, SEED)

Postemergence Application

Apply as a postemergence broadcast spray using sufficient spray volume to ensure thorough coverage of treated plants, but no less than 15 gpa for ground spray equipment or 2 to 5 gpa for aircraft equipment. Control may be reduced at low spray volumes under high temperature and wind conditions. Lorsban Advanced may be tank mixed with glyphosate products, such as Duramax herbicide or Durango DMA herbicide, when application is to be made to glyphosate-tolerant corn.

Chemigation: Lorsban Advanced may be broadcast applied postemergence through sprinkler irrigation systems at specified application rates to control listed foliar pests. For best results, tank mix Lorsban Advanced with 2 pints of non-emulsifiable oil. See Chemigation Application section.

Pest-Specific Use Directions:

Billbug, chinch bug, or flea beetle: For best control, ground apply in a minimum spray volume of 20 to 40 gpa at 40 psi. If corn is less than 6 inches tall, apply in a 9- to 12-inch wide band over the row. For corn more than 6 inches tall, apply using drop nozzles directed to the base of the plant. Do not reduce the application rate for banded or directed applications. Concentrate the full labeled dosage rate in the treated zone. When chinch bugs continue to immigrate to corn over a prolonged period or under extreme pest pressure, a second application may be needed.

Corn rootworm larvae: For postemergence control, apply at cultivation. Direct the spray to both sides of the row at the base of the plants just ahead of the cultivator shovels. Cover the insecticide with soil around the brace roots. A cultivation application of Lorsban Advanced may be made in addition to an at-planting application of Lorsban 15G.

Lorsban Advanced may also be applied through sprinkler irrigation systems at the rate of 2 pints per acre to control corn rootworm larvae. Time application to coincide with the appearance of the second instar larvae. Apply with enough water to wet the root zone to the depth control needed. If soils are wet, allow enough soil drying to occur such that an application using a minimum amount of water will not produce surface runoff. See Chemigation Application section for application instructions.

Do not use Lorsban Advanced in combination with a burndown herbicide for control of common stalk borer. For common stalk borer control, treat approximately 11 days after application of glyphosate or after burndown with paraquat herbicide is complete (3 to 5 days).

Specific Use Restrictions:

- Preharvest Interval: Do not apply within 21 days before harvest of grain, ears, forage or fodder.
- Do not make more than three applications of Lorsban Advanced or any product containing chlorpyrifos per season, including the maximum allowed of two granular applications, at the 1 lb ai chlorpyrifos rate.
- Do not make a second application of Lorsban Advanced or other product containing chlorpyrifos within 10 days of the first application.
- If more than 1 lb ai granular chlorpyrifos per acre is applied at-plant (for a maximum of 1.3 lb ai per acre per season), only one additional application of a liquid product containing chlorpyrifos at 1 lb ai per acre is allowed per season, for a total of 2.3 lb ai chlorpyrifos per acre per season.
- Do not apply in tank mixes with Steadfast or Lightning herbicides.
- Do not aerially apply this product in Mississippi.

Method

[Broadcast/Foliar Air](#)

[Broadcast/Foliar Ground](#)

Pre-Harvest Interval

21 days

Rates

[field rates 0](#)

•

Restricted Entry Interval

24 hours

Timings

[Postemergence \(Crop\)](#)