

COMMERCIAL FLORIST - OVERHEAD WATERING SYSTEMS

General Information

GENERAL DISINFECTION PERFORMANCE

This product kills the following bacteria in 10 minutes at 1.5 oz. per 5 gal. of water (469 ppm active) and 5% soil on hard, non-porous surfaces:

Salmonella enterica

Staphylococcus aureus

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Please read entire label and use strictly in accordance with precautionary statements and directions.

Rinse hard, non-porous food contact surfaces with potable water after application of product.

Limitations, Restrictions, and Exceptions

COMMERCIAL FLORIST

DO NOT apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirement specific to your State and Tribe, consult the State/Tribal agency responsible for pesticide regulation.

OVERHEAD WATERING SYSTEMS: During the last 2-3 minutes of watering use 0.5 oz. of this product per 20 gal. of water (or equivalent use dilution) as a control program. Due to variations of growing conditions, use every second to fourth watering. Will control algae in nozzles.

Method

[Water System Treatment](#)

Rates

[field rates 0](#)

-

Restricted Entry Interval

48 hours

Timings

[Every second to fourth watering.](#)