

SOIL-INJECTED OR SOIL-INCORPORATED APPLICATIONS - GRAPES - SOIL SPRAY

General Information

FORESTRY APPLICATION INSTRUCTIONS

Consult with recognized forest pest management authorities or an AEF Global Inc. representative regarding appropriate timing and method of application. The timing and number of applications required for effective forest protection will depend upon the target pest, target foliage development, pest pressure and larval activity. Correspond applications with sufficient foliage development (shoot elongation or leaf expansion) to ensure maximum spray deposit. To the extent possible, apply treatments after egg hatch has been completed to assure that the maximum number of larvae is present during the treatment period. In conditions where egg hatch is extended and/or larval development is significantly advanced and/or larval populations are high, use of higher rates and/or additional applications, 3-10 days apart, is recommended. See dose rate table for pest-specific application instructions.

Ground Application:

Dilute required amount of Bioprotec PLUS with water only to the volume required to provide thorough coverage. Do not wet foliage to the point of excessive runoff.

For mist blower applications, add Bioprotec PLUS to water at a ratio of 1:20 to 1:50. Use a maximum of 10.7 gallons of spray mixture per acre.

For high volume hydraulic sprayers, add Bioprotec PLUS to water at a ratio of 1:50 to 1:500. Use a maximum of 107 gallons of spray mixture per acre.

Aerial Application:

Apply only by fixed-wing or rotary aircraft equipment that has been functionally and operationally calibrated for the atmospheric conditions of the area and the application rates and conditions of this label. Label rates, conditions and precautions are product specific.

Apply only at the rate specified for aerial application on this label. Where no rate for aerial application appears for the specific use, this product cannot be applied by any type of aerial equipment. Ensure uniform application by using appropriate

marking devices and/or electronic guidance equipment.

Undiluted applications are highly recommended for most labeled pests. To improve coverage, Bioprotec PLUS may be diluted with water and applied at the rates indicated in the dose rates table. Total volume of spray material to be applied per acre depends upon the target pest, target foliage, weather, spray equipment and droplet size. Best results are expected when Bioprotec PLUS is applied undiluted to dry foliage with well calibrated aircraft and appropriate spray systems capable of delivering droplets in the range of 500-170 U.S. mesh (30-80 microns) for coniferous foliage and 270-100 U.S. mesh (50-150 microns) for broadleaf foliage.

FOR AGRICULTURAL USES:

For use on trees or shrubs being grown for sale or other commercial use, such as for commercial seed production or research purposes.

- Use the higher rates on advanced larval stages or under high density larval populations.

Limitations, Restrictions, and Exceptions

SOIL-INJECTED OR SOIL-INCORPORATED APPLICATIONS

The restricted-entry interval (REI) for soil-injected or soil-incorporated applications is 0 hours.

GRAPE

Application Directions

- Soil spray (broadcast, band or drip): Apply in the spring before the plants start growing in sufficient water to provide uniform coverage. Two additional applications may be made to coincide with periods most favorable for root rot development.
- Soil spray (broadcast or band): For banded applications, use a 3-ft band at the base of the plants. Calculate the correct amount based on 3.6 pt/A using the instructions in the Application Procedures section.

Use Restrictions:

1. Do not exceed the equivalent of 5.4 lb ai/A per crop of soil-applied mefenoxam-containing products and 0.4 lb ai/A per crop of foliar-applied mefenoxam-containing products.

Method

[Broadcast/Foliar Air](#)

[Broadcast/Foliar Ground](#)

[Band application](#)

[Drip](#)

[Soil Spray](#)

Pre-Harvest Interval

60 days

Rates

[field_rates 0](#)

[field_rates 1](#)

[field_rates 2](#)

•

Restricted Entry Interval

0 hours

Timings

[In the spring before the plants start growing](#)