

FOOD/FEED HANDLING ESTABLISHMENTS - NON-FOOD/FEED AREAS

General Information

Directions For Use

It is a violation of federal law to use this product in a manner inconsistent with its labeling.

DO NOT apply this product while bees are foraging. DO NOT apply this product to plants that are flowering. Only apply after all flower petals have fallen off.

Use Restrictions

- DO NOT apply within aircraft cabins.
- DO NOT use in outdoor residential misting systems.
- DO NOT apply to humans or their clothing or to mattresses. DO NOT allow children or pets to contact treated surfaces until spray has dried.
- DO NOT apply to household pets. Cover fish tanks prior to application near tanks.
- DO NOT apply this product in patient rooms while occupied.
- DO NOT apply to classrooms when in use.
- When applying in residential or commercial structures, DO NOT contaminate food, feed stuffs or water supply. DO NOT contaminate food preparation surfaces, kitchen utensils, dishes or food/feed storage containers. Cover any food/feed contact surfaces and cooking utensils in the treatment area before treatment or thoroughly clean after treatment and before using.
- Apply to the point just prior to runoff. DO NOT allow applications of product to drip or run from surfaces.
- When applying to structural areas in and around livestock housing, remove animals or treat in areas where animals are not present. DO NOT apply to animals. DO NOT allow insecticide to drift onto animals. DO NOT contaminate milk, food or drinking water.

Product Information

This product is a water-soluble granule containing the non-repellent active ingredient dinotefuran. Apply indoors and/or outdoors via crack and crevice, spot or surface application for the listed pests and use sites.

Mixing Instructions

Thoroughly clean spray equipment before mixing and application of this product.

Add approximately 1/2 of the water to the spray tank. Add 10 to 30 grams of product to water per 1,000 square foot area being treated (1 Tablespoon = 10 grams), then complete the filling of the spray tank. Agitate the tank until the product is completely dissolved. Reagitate finished dilution if application is interrupted for an extended period of time.

Void/Crack & Crevice Application

Apply into cracks, crevices and voids with equipment designed for such applications. Follow the manufacturer's recommendations for proper equipment set-up and operating pressures. Place the tip of the applicator at or into the crack, crevice or void to be treated. Apply product in short bursts while keeping tip in position. Allow the air stream to push the finished dilution into voids and/or cracks and crevices. Apply in such a manner to avoid dripping and runoff.

Limitations, Restrictions, and Exceptions

Food/Feed Handling Establishments

Food/Feed handling establishments are places other than private residences in which food is held, processed, prepared or served, including those operating under the Federal meat, poultry, shell egg grading and egg products inspection programs. APPLICATIONS OF THIS PRODUCT IN FOOD/FEED AREAS OF FOOD/FEED HANDLING ESTABLISHMENTS MAY BE MADE AS A CRACK & CREVICE, VOID, SPOT OR GENERAL TREATMENT. General surface application may be used only when the facility is not in operation provided exposed food/feed has been covered or removed from the area prior to application.

Non-food/feed areas include areas such as garbage rooms, lavatories, floor drains (to sewers), entries and vestibules, offices, locker rooms, machine rooms, boiler rooms, garages, mop closets and storage areas (after packaging, canning or

bottling).

Avoid contamination of food/feed or food/feed contact surfaces. Remove or cover food/feed, dishes, utensils, food processing equipment and food preparation surfaces in the treatment area or wash them before use. Apply as a Crack & Crevice, void, spot or general treatment to surfaces such as baseboards, under elements of construction, stainless steel equipment, shelving, machinery, storage areas, pallets, tables, chairs and other areas where listed pests may be harboring, traveling, breeding or entering the structure. Maximum use rate is 0.05 grams of active ingredient per linear foot. Reapplications may be made at 3 day intervals when applying as a Crack & Crevice, void or spot treatment, or at 6 day intervals when applying as a general surface treatment. This product can also be used as a preventative treatment for areas likely to have future infestations.

Method

[Spot treatment](#)

Rates

[field rates 0](#)

-