

INDOOR APPLICATIONS - KILL HOUSE FLIES

General Information

DO NOT apply this product while bees are foraging. DO NOT apply this product to plants that are flowering. Only apply after all flower petals have fallen off.

USE RESTRICTIONS:

- DO NOT apply within aircraft cabins.
- DO NOT use in outdoor residential misting systems.
- DO NOT apply to humans or their clothing or to mattresses. DO NOT allow children or pets to contact treated surfaces until spray has dried.
- DO NOT apply to household pets. Cover fish tanks prior to application near tanks.
- DO NOT apply this product in patient rooms while occupied.
- DO NOT apply to classrooms when in use.
- When applying in residential or commercial structures, DO NOT contaminate food, feed stuffs or water supply. DO NOT contaminate food preparation surfaces, kitchen utensils, dishes or food/feed storage containers. Cover any food/feed contact surfaces and cooking utensils, in the treatment area, before treatment or thoroughly clean after treatment and before using.
- Apply to the point just prior to runoff. DO NOT allow application to drip or run from surfaces.

PRODUCT INFORMATION

This product is a water-soluble granule containing the nonrepellent active ingredient Dinotefuran. Apply indoors and/or outdoors via crack and crevice, spot or surface application for the listed pests and use sites.

Refer to specific pest categories below for the proper use rate. One gallon of finished dilution treats 1,000 sq ft. Depending on the surface, more or less water may be used to properly treat areas with different application apparatus, but the proper amount of active ingredient should be used per square footage.

Tank Mixing: This product may be tank mixed with other water based insecticides including, but not limited to insect growth regulators (IGR), quick knockdown products and residuals. Before mixing test to see if products are compatible. When

tank mixing, follow the more restrictive label use directions of the mixed products.

Limitations, Restrictions, and Exceptions

TO KILL HOUSE FLIES INDOORS AND OUTDOORS

Indoor Applications

Using the 0.1% to 0.2% application rate, make spot applications on surfaces where flies frequently roost such as door and window frames, screens, trash cans, plants, interiorscapes, ceilings and upper wall surfaces.

Method

[Spot treatment](#)

Rates

[field rates 0](#)

-

Timings

[N.A.](#)