

# **HARVEST AID APPLICATIONS ON COTTON (TANK MIXES WITH PHOSPHATE OR CHLORATE DEFOLIANTS) - TEXAS**

Limitations, Restrictions, and Exceptions

HARVEST AID APPLICATIONS ON COTTON IN TEXAS

(except in the counties of Starr, Hidalgo, Willacy, and Cameron).

TANKMIXES WITH PHOSPHATE OR CHLORATE DEFOLIANTS

Rate: 8.1 – 16.1 fl oz. alone or tank mixed with 1 1/2 - 2 gal. chlorate, or 1 1/2 - 2 pts. phosphate

Precautions, Restrictions and Comments

- Do not harvest within 7 days of application when used with phosphate or chlorate defoliants.
- In desert cotton areas or on rank vigorous cotton, use the 16.1 fl oz rate of Gramoxone SL 2.0
- Development of immature bolls will be inhibited.
- Follow Harvest Aid Use Precautions listed on the Federal label.

Specific Crop Use Directions

Three multiple broadcast harvest aid applications of Gramoxone SL 2.0 may be used. Use a minimum total spray volume per acre of 10 gallons by ground, and 5 gallons by air.

Refer to the Gramoxone SL 2.0 federal label for additional cotton recommended uses and nonionic surfactant recommendations.

Additional Precautions, Restrictions and Comments:

- It is advisable, because of extremes in environmental and plant conditions, to apply the range of rates on a small block of cotton to determine the rate that best fits your needs.
- Do not pasture lactating dairy animals.
- Do not pasture livestock in treated fields or feed treated foliage.
- When combined with phosphate defoliants, observe livestock cautions listed on

the phosphate label.

- Do not harvest within 7 days of application when used with phosphate or chlorate defoliants.
- Do not apply this product through any type of irrigation system.

Method

[Broadcast/Foliar Air](#)

[Broadcast/Foliar Ground](#)

Pre-Harvest Interval

7 days

Rates

[field rates 0](#)

- 

Restricted Entry Interval

24 hours

Timings

[When 75% or more of the bolls are open and remaining bolls to be harvested are mature.](#)