

HARVEST AID APPLICATIONS ON COTTON (TANK MIXES FOR BOLL OPENING AND DEFOLIATION OF SOUTHERN COTTON) - TEXAS

Limitations, Restrictions, and Exceptions

HARVEST AID APPLICATIONS ON COTTON IN TEXAS

(except in the counties of Starr, Hidalgo, Willacy, and Cameron).

TANKMIXES FOR BOLL OPENING AND DEFOLIATION OF SOUTHERN COTTON

Precautions, Restrictions and Comments

- To aid in defoliation and opening of mature bolls, Gramoxone SL 2.0 may be tank mixed with the following products:
DEF defoliant, DROPP defoliant, PREP PGR, FOLEX defoliant
- Apply when 60% or more of the bolls are open and the remaining bolls to be harvested are mature.
- Development of immature bolls will be inhibited.
- Refer to tank mix product label for rate, directions, limitations and caution.

Specific Crop Use Directions

Three multiple broadcast harvest aid applications of Gramoxone SL 2.0 may be used. Use a minimum total spray volume per acre of 10 gallons by ground, and 5 gallons by air.

Refer to the Gramoxone SL 2.0 federal label for additional cotton recommended uses and nonionic surfactant recommendations.

Additional Precautions, Restrictions and Comments:

- It is advisable, because of extremes in environmental and plant conditions, to apply the range of rates on a small block of cotton to determine the rate that best fits your needs.
- Do not pasture lactating dairy animals.
- Do not pasture livestock in treated fields or feed treated foliage.
- When combined with phosphate defoliants, observe livestock cautions listed on the phosphate label.

- Do not harvest within 7 days of application when used with phosphate or chlorate defoliant.
- Do not apply this product through any type of irrigation system.

Method

[Broadcast/Foliar Air](#)

[Broadcast/Foliar Ground](#)

Pre-Harvest Interval

7 days

Rates

[field_rates 0](#)

•

Restricted Entry Interval

24 hours

Timings

[When 60% or more of the bolls are open and the remaining bolls to be harvested are mature.](#)