

HARVEST AID APPLICATIONS ON COTTON (DESSICATION OF REGROWTH) - TEXAS

Limitations, Restrictions, and Exceptions

HARVEST AID APPLICATIONS ON COTTON IN TEXAS

(except in the counties of Starr, Hidalgo, Willacy, and Cameron).

DESSICATION OF REGROWTH

Precautions, Restrictions and Comments

- Use to desiccate regrowth after defoliation or desiccation.
- Regrowth is difficult to control, therefore, thorough coverage with the full recommended rate is necessary.
- Control is dependent upon growing condition and desiccation of small new growth may not always be complete.
- Use higher rate if regrowth is excessive.

Specific Crop Use Directions

Three multiple broadcast harvest aid applications of Gramoxone SL 2.0 may be used. Use a minimum total spray volume per acre of 10 gallons by ground, and 5 gallons by air.

Refer to the Gramoxone SL 2.0 federal label for additional cotton recommended uses and nonionic surfactant recommendations.

Additional Precautions, Restrictions and Comments:

- It is advisable, because of extremes in environmental and plant conditions, to apply the range of rates on a small block of cotton to determine the rate that best fits your needs.
- Do not pasture lactating dairy animals.
- Do not pasture livestock in treated fields or feed treated foliage.
- When combined with phosphate defoliant, observe livestock cautions listed on the phosphate label.
- Do not harvest within 7 days of application when used with phosphate or chlorate defoliant.

- Do not apply this product through any type of irrigation system.

Method

[Broadcast/Foliar Air](#)

[Broadcast/Foliar Ground](#)

Pre-Harvest Interval

7 days

Rates

[field_rates 0](#)

•

Restricted Entry Interval

24 hours

Timings

[N.A.](#)