

VEGETABLE AND SMALL FRUIT CROPS - GLOBE ARTICHOKE - SOIL

General Information

APPLICATION DIRECTIONS

Soil Application:

Direct applications of CONDOR into the seed or root-zone of crop. Failure to place CONDOR into root-zone may result in loss of control or delay in onset of activity. Apply CONDOR by ground application or chemigation application. For seedling flats or trays, only apply with broadcast, foliar applications or where product is intended to be washed from foliage to soil prior to drying on foliage.

Optimum activity of CONDOR results from applications to the root-zone of plants to be protected. The earlier CONDOR is available to a developing plant, the earlier the protection begins. CONDOR is continuously taken into the roots over a long period of time and the systemic nature of CONDOR allows movement from roots through the xylem tissue to all vegetative parts of the plant. This results in extended residual activity of CONDOR, the control of insects and the prevention and/or reduction of virus transmission or symptom expression, and plant health benefits. The rate of CONDOR applied affects the length of the plant protection period. Use the higher listed rates when infestations occur later in crop development, or where pest pressure is continuous. CONDOR will generally not control insects infesting flowers, blooms or fruit. Additional crop protection may be required for insects feeding in, or on these plant parts and for insects not listed in the crop-specific, pests controlled sections of this label. Additional, specific CONDOR application rates are also provided in the crop-specific sections of this label.

RESTRICTIONS:

- Do not apply with aerial application equipment
- Do not apply more than 0.5 lb active ingredient per acre, per year regardless of formulation or method of application, unless specified within a crop-specific section for a given crop.

Foliar Application:

Do not apply CONDOR in enclosed structures such as greenhouses or planthouses. Apply CONDOR as a directed or broadcast foliar spray. Thorough coverage of foliage is necessary without runoff for optimum insecticidal efficacy.

Use adequate spray volumes, properly calibrated application equipment, and spray adjuvant if necessary to obtain thorough coverage. Failure to provide adequate coverage and retention of CONDOR on leaves and fruit may result in loss of insect control or delay in onset of activity. CONDOR may be applied with properly calibrated ground or aerial application equipment. Minimum specified spray volumes unless otherwise specified on crop specific application sections are 10 gallons/acre by ground application and 5 gallons/acre through aerial equipment. CONDOR may also be applied by overhead chemigation (see additional Chemigation Directions for Use section below) if allowed in crop specific Application section. CONDOR use on crops grown for production of true seed intended for private or commercial planting is not permitted unless specifically approved under state-specific 24(c) Special Local Needs labeling. Additional information on CONDOR uses for these crops and other questions may be obtained from the Cooperative Extension Service, PCAs, consultants, or local Altitude Crop Innovations, LLC representatives.

RESTRICTIONS:

- Do not apply to plants grown in non-soil medias such as perlite, vermiculite, rock wool or other soil-less media, or plants growing hydroponically.
- Do not apply more than 0.5 lb active ingredient per acre, per year regardless of formulation or method of application, unless specified within a crop-specific section for a given crop.

CHEMIGATION

Refer to Directions For Use section before proceeding with chemigation application. For Soil Application: Chemigation applications of CONDOR may only be made to crops through chemigation systems as specified in crop-specific Application Instructions section and only through low-pressure systems unless specified for a given crop. DO NOT apply CONDOR through any other type of irrigation system. For Foliar Application: Chemigation applications of CONDOR may be made to crops through overhead sprinkler chemigation if specified in crop specific instruction sections. DO NOT apply CONDOR through any other type of irrigation system. Make foliar chemigation applications of CONDOR as concentrated as possible. Retention of CONDOR on target site of insect infestation is necessary for optimum activity. DO NOT chemigate CONDOR in water volumes exceeding 0.10 inch/Acre.

Water Volume

CONDOR chemigation applications should be made as concentrated as possible. Retention of CONDOR on target site of insect infestation is necessary for optimum

activity. Do not chemigate CONDOR in water volumes exceeding 0.10 inch/acre.

Limitations, Restrictions, and Exceptions

Restrictions:

- Maximum CONDOR allowed per year when making soil applications: 16 fluid ounces per acre (0.5 lb AI/Acre).

Applications: Apply specified dosage in one of the following methods:

1. Chemigation into root-zone through low-pressure drip, trickle, micro-sprinkler or equivalent equipment;
2. In-furrow spray at planting directed on or below seed.

Method

[In Furrow Spray](#)

Pre-Harvest Interval

7 days

Rates

[field_rates 0](#)

•

Restricted Entry Interval

12 hours

Exception: If the product is soil-injected or soil-incorporated, the Worker Protection Standard, under certain circumstances, allows workers to enter the treated area if there will be no contact with anything that has been treated.

Timings

[At-Plant](#)