

TURFGRASS RENOVATION, SEED OR SOD PRODUCTION - PERENNIAL WEEDS

General Information

Product Information

This product is a broad spectrum, systemic, postemergent herbicide with no soil residual activity. It is intended for control of annual and perennial weeds and woody plants and brush. It is formulated as a water soluble liquid.

Time to Symptoms: The active ingredient in this product moves through the plant from the point of foliage contact to and into the root system. Visible effects are a gradual wilting and yellowing of the plant that advances to complete browning of above ground growth and deterioration of underground plant parts. Visible effects on most annual weeds occur within two to four days, but on most perennial weeds visible effects may not occur for seven days or more. Extremely cool or cloudy weather following treatment may slow the activity of this product and delay development of visual symptoms.

Stage of Weeds: Annual weeds are easiest to control when they are small. Best control of most perennial weeds is obtained when treatment is made at late growth stages approaching maturity. Refer to the annual, perennial and woody brush and trees rate tables for specific weeds. Always use the higher rate within the rate range for heavy or dense weed growth or when weeds are growing in an undisturbed (non-cultivated) area. When treating weeds with disease or insect damage, weeds heavily covered with dust, or weeds under poor growing conditions, reduced weed control may result.

Cultural Considerations: Reduced control may result when applications are made to annual or perennial weeds that have been mowed, grazed, or cut, and have not been allowed to regrow to the specified stage for treatment.

Rainfastness: Heavy rainfall soon after application may wash off this product from the foliage and a repeat application up to the labeled rate may be required for adequate control.

Spray Coverage: For best results, spray coverage should be uniform and complete.

Mode of Action: The active ingredient in this product inhibits an enzyme. This enzyme is found only in plants and microorganisms that are essential to forming specific amino acids.

No Soil Activity: Weeds must be emerged at the time of application to be controlled by this product. Weeds germinating from seed after application will not be controlled. Unemerged plants arising from unattached underground rhizomes or rootstocks of perennials will not be affected by the herbicide and will continue to grow.

Biological Degradation: Degradation of this product is primarily a biological process carried out by soil microbes.

Maximum Application Rates: The maximum application rates specified in this label are given in units of volume, either fluid ounces, pints or quarts, of this product per acre. The maximum allowed application rates apply to this product combined with the use of any and all other glyphosate- or sulfosate-containing herbicides, either applied separately or in a tank mix, on the basis of total pounds of glyphosate (acid equivalents) per acre. If more than one glyphosate- or sulfosate-containing product is applied to the same site within the same year, ensure that the total of pounds acid equivalent glyphosate does not exceed the maximum allowed.

Do not apply more than 8 quarts of this product (8 lb glyphosate acid) per acre per year for all use sites listed on this label.

IMPORTANT: When using this product, unless otherwise specified, mix with a surfactant, such as a nonionic surfactant containing 80% or greater active ingredient. For conifer release (pine release) use only surfactants that are approved for conifer release and specified on the surfactant label as safe for use in conifer release (pine release). Use of this product without surfactant will result in reduced herbicide performance. Ammonium sulfate, drift control additives, or dyes and colorants may be used. See Mixing Directions and the surfactant manufacturer's label for more information.

Grazing Restrictions: This product may be used to treat undesirable vegetation in utility rights-of-way that pass through pastures, rangeland, and forestry sites that

are being grazed. For tank mix applications, comply with all restrictions appearing on the tank mix product label.

Except for lactating dairy animals there are no grazing restrictions following the labeled applications of this product.

For lactating dairy animals there are no grazing restrictions for the following labeled applications of this product:

- Where the spray can be directed onto undesirable woody brush and trees, including in handgun spray to wet or low volume directed spray treatments.
- For tree injection of frill applications and for cut stump treatments.

For broadcast applications, observe the following restrictions for lactating dairy animals:

- For application rates between 4.5 and 7.5 quarts per acre, no more than 15 percent of the available grazing area may be treated.
- For application rates less than 4.5 quarts per acre, no more than 25 percent of the available grazing area may be treated.

These restrictions do not apply to pastures, rangeland or forestry sites outside of utility rights-of-way.

Herbicide Resistance Management

Glyphosate, the active ingredient in this product, is a group 9 herbicide (inhibitor of EPSP synthase). Some naturally occurring weed biotypes that are tolerant (resistant) to glyphosate may exist due to genetic variability in a weed population. Where resistant biotypes exist, the repeated use of herbicides with the same mode of action can lead to the selection for resistant weeds. Certain agronomic practices reduce the likelihood that resistant weed populations will develop, and can be utilized to manage weed resistance once it occurs.

To delay the selection for glyphosate resistant weeds, use the following practices:

- Scout fields before and after application to detect weed escapes or shifts in weed species.
- Start with a clean field by applying a burndown herbicide or by tillage.
- Control weeds early when they are small.
- Add other herbicides, including a selective and/or a residual herbicide, and cultural practices, including tillage or crop rotation, where appropriate.

- Use the application rate for the most difficult to control weed in the field. Do not tank mix with other herbicides that reduce this product's efficacy through antagonism or with ones that encourage application rates of this product below those specified on this label.
- Control weed escapes and prevent weeds from setting seeds.
- In situations where resistant weeds are a problem, before moving from one site to another, clean equipment to minimize the spread of weed seeds or plant parts.
- Use new commercial seed that is as free of weed seed as possible.
- Report any incidence of repeated non-performance of this product against a particular weed species to the local retailer, county extension agent, or Dow AgroSciences representative.

The following good agronomic practices are recommended to reduce the spread of confirmed glyphosate-resistant biotypes:

- Tank mix this product or apply it sequentially with an appropriately labeled herbicide with a different mode of action to achieve control if a naturally occurring resistant biotype is present in the site.
- Cultural and mechanical control practices, including crop rotation or tillage, may also be used.
- To control weed escapes, including resistant biotypes, before they set seed, scout treated sites after applying this product.
- Thoroughly clean equipment before leaving any site known to contain resistant biotypes.

Because the presence of glyphosate resistance in weed populations is difficult to detect prior to use, Dow AgroSciences accepts no liability for any losses that may result from the failure of this product to control glyphosate-resistant weeds.

Attention

Avoid contact of herbicide with foliage, green stems, exposed non-woody roots or fruit of crops, desirable plants and trees, because severe injury or destruction may result.

AVOID DRIFT. Use extreme care when applying this product to prevent injury to desirable plants and crops.

Do not allow the herbicide solution to mist, drip, drift or splash onto desirable vegetation since minute quantities of this product can cause severe damage or

destruction to the crop, plants or other areas on which treatment was not intended. The likelihood of injury occurring from the use of this product increases when winds are gusty, as wind velocity increases, when wind direction is constantly changing, or when there are other meteorological conditions that favor spray drift. When spraying, avoid combinations of pressure and nozzle type that will result in splatter or fine particles (mist) which are likely to drift. Avoid applying at excessive speed or pressure.

NOTE: Use of this product in any manner not consistent with this label may result in injury to persons, animals or crops, or other unintended consequences. Keep container closed to prevent spills and contamination.

Application Equipment and Application Methods

Chemigation: Do not apply this product through any type of irrigation system.

Apply spray solutions in properly maintained and calibrated equipment capable of delivering desired volumes.

This product may be applied with the following application equipment and application methods.

Aerial Application

Equipment: Fixed wing and helicopter

Do not apply this product using aerial spray equipment except under conditions as specified within this label.

Avoid drift. Do not apply when winds are gusty or under any other condition which favors drift. Drift may cause damage to any vegetation contacted to which treatment is not intended. To prevent injury to adjacent desirable vegetation, maintain appropriate buffer zones.

Do not directly apply to any body of water.

Use the specified rates of this herbicide in 3 to 25 gallons of water per acre unless otherwise specified on this label. Refer to the specific use directions of this label for volumes and application rates.

Coarse sprays are less likely to drift; therefore, do not use nozzles or nozzle

configurations that dispense spray as fine spray droplets. Do not angle nozzles forward into the airstream and do not increase spray volume by increasing nozzle pressure. A drift control additive may be used. When a drift control additive is used, carefully read and observe the precautionary statements and all other information specified on the additive label.

Ensure uniform application. To avoid streaked, uneven or overlapped application, use appropriate marking devices.

Aerial Application Restrictions in California Only

AVOID DRIFT: Do not apply when winds are gusty or under any other condition which favors drift. Drift may cause damage to any vegetation contacted to which treatment is not intended. To prevent injury to adjacent desirable vegetation, appropriate buffer zones must be maintained.

Do not aerially apply this product in a tank mix with dicamba in California.

Make aerial applications with helicopter only. To ensure uniform application, avoid streaking, uneven, or overlapped application, and use appropriate marking devices.

Use the following guidelines when aerial applications are made near crops or desirable perennial vegetation after budbreak and before total leaf drop, and/or near other desirable vegetation or annual crops:

- Do not apply this product using aerial equipment in residential areas.
- Do not apply within 100 feet of all desirable vegetation or crop(s).
- If wind up to 5 miles per hour is blowing toward desirable vegetation or crop(s), do not apply within 500 feet of the desirable vegetation or crop(s).
- Winds blowing from 5 to 10 miles per hour toward desirable vegetation or crop(s) may require buffer zones in excess of the 500-foot minimum buffer.
- Do not apply when winds are in excess of 10 miles per hour or when inversion conditions exist.

Use only coarse sprays to minimize drift. Do not use nozzles or nozzle configurations that dispense spray as fine spray droplets. Do not angle nozzles forward into the airstream and do not increase spray volume by increasing nozzle pressure above the manufacturer's directions.

Thoroughly wash aircraft, especially landing gear, after each day of spraying to

remove residues of this product accumulated during spraying or from spills. Prolonged exposure of this product to uncoated steel surfaces may result in corrosion and possible failure of the part. Landing gear is most susceptible. The maintenance of an organic coating (paint) which meets aerospace specification MIL-C-38413 may prevent corrosion.

ADDITIONAL LIMITATIONS FOR AERIAL APPLICATION IN FRESNO COUNTY, CALIFORNIA ONLY

Always read and follow the label directions and precautionary statements for all products used in the aerial application.

The following information applies only from February 15 through March 31 within the following boundaries of Fresno County, California:

North: Fresno County line

South: Fresno County line

East: State Highway 99 West

Observe the following directions to minimize off-site movement during aerial application of this product. Minimization of off-site movement is the responsibility of the grower, Pest Control Advisor and aerial applicator.

Written Directions

Written directions MUST be submitted by or on behalf of the applicator to the Fresno County Agricultural Commissioner 24 hours prior to the application. These written directions MUST state the proximity of surrounding crops and that conditions of each manufacturer's product label and this label have been satisfied.

Aerial Applicator Training and Equipment

Aerial application of this product is limited to pilots who have successfully completed a Fresno County Agricultural Commissioner and California Department of Pesticide Regulation approved training program for aerial application of herbicides. All aircraft must be inspected, critiqued in flight and certified at a Fresno County Agricultural Commissioner approved fly-in. Test and calibrate spray equipment at intervals sufficient to insure that proper rates of herbicides and adjuvants are being applied during commercial use. Applicator must document such calibrations and testing. Demonstration of performance at Fresno County Agricultural Commissioner approved fly-ins constitutes such documentation, or other written records showing calculations and measurements of flight and spray parameters acceptable to the

Fresno County Agricultural Commissioner.

Applications at Night – Do not apply this product by air earlier than 30 minutes prior to sunrise and/or later than 30 minutes after sunset without prior permission from the Fresno County Agricultural Commissioner.

To report known or suspected misuse of this product, call 1-800-332-3111.

For additional information on the proper aerial application of this product in Fresno County, call 916-784-1718.

Aquatic and Noncrop Sites

When this product is applied under the conditions described, it controls or partially controls the labeled weeds growing in the following industrial, recreational, and public areas or other similar sites.

Aquatic sites includes all bodies of fresh and brackish water that may be flowing, nonflowing, or transient including lakes, rivers, streams, ponds, seeps, irrigation and drainage ditches, canals, reservoirs, estuaries and similar sites.

If aquatic sites are present in the noncrop area and are part of the intended treatment, read and observe the following directions:

- This product does not control plants that are completely submerged or have a majority of their foliage under water.
- There is no restriction on the use of treated water for irrigation, recreation, or domestic purposes.
- Consult local and state fish and game agency and water control authorities before applying this product to public water. Permits may be required to treat such water.
- To make aquatic applications around and within 1/2 mile of active potable water intakes, the water intake must be turned off for a minimum period of 48 hours after the application. The water intake may be turned on prior to 48 hours if the glyphosate level in the intake water is below 0.7 parts per million as determined by laboratory analysis. These aquatic applications may be made only in those cases where there are alternative water sources or holding ponds that would permit the turning off of an active potable water intake for a minimum period of 48 hours after the application.

Restrictions:

- Do not apply this product within 1/2 mile upstream of an active potable water

intake in flowing water (i.e., river stream, etc.), or within 1/2 mile of an active potable water intake in a standing body of water, such as a lake, pond, or reservoir.

Ground Application

Equipment: Boom or boomless systems, pull-type sprayer, floaters, pick-up sprayers, spray coupes and other ground broadcast equipment.

Use the specified rates of this product in 3 to 40 gallons of water per acre as a broadcast spray unless otherwise specified on this label. As density of weeds increases, increase the spray volume within the rate range to ensure complete coverage. Carefully select proper nozzles to avoid spraying a fine mist. For best results with ground application equipment, use flat fan nozzles. Check for even distribution of spray droplets.

Hand-Held and High-Volume Including Backpack Application

Equipment: Knapsack and backpack sprayers, pump up pressure sprayers, handguns, hand wands, mistblowers, lances, and other hand-held and motorized spray equipment used to direct the spray onto weed foliage. Note: This product is not registered in Arizona or California for use in mistblowers.

Apply to foliage of vegetation to be controlled. Do not spray to the point of runoff for applications made on a spray to wet basis. Use coarse sprays only. For best results, cover the top half of the plant and at least half of the total foliage. To ensure adequate spray coverage, spray both sides of large or tall woody brush and trees, when foliage is thick and dense, or where there are multiple sprouts.

High Volume Sprays: Prepare a 3/4 to 2 percent solution of this product in water, add a nonionic surfactant and apply to foliage of vegetation to be controlled. For specific rates of application and instructions for control of various annual and perennial weeds, see the Weeds Controlled section.

Make applications on a spray to wet basis with uniform and complete spray coverage. Do not spray to point of runoff.

Low Volume Directed Sprays: This product may be used as a 5 to 10 percent solution in low volume directed sprays for spot treatment of trees and brush. This treatment method is most effective in areas where there is a low density of undesirable trees or brush. If a straight stream nozzle is used, start the application at the top of the targeted vegetation and spray from top to bottom in a lateral

zigzag motion. Ensure that at least 50 percent of the leaves are contacted by the spray solution. For flat fan and cone nozzles and with hand-directed mist blowers, mist the application over the foliage of the targeted vegetation. Treat small, open-branched trees only from one side. If the foliage is thick or there are multiple root sprouts, apply from several sides to ensure adequate spray coverage. Prepare the desired volume of spray solution by mixing the amount of this product in water as shown in the following table.

Spray Solution:

For best results when using knapsack sprayers, mix the specified amount of product with water in a larger container. Fill the knapsack sprayer with the solution and add the correct amount of surfactant.

Selective Equipment

Equipment: Recirculating sprayers, shielded and hooded sprayers, wiper applicators and sponge bars.

Do not contact desirable vegetation with herbicide. Droplets, mist, foam, or splatter of the herbicide settling on desirable vegetation is likely to result in discoloration, stunting or destruction.

Better results are obtained when more of the weed is exposed to the herbicide solution. Weeds not contacted by the herbicide solution will not be affected. This may occur in dense clumps, severe infestations, or when the height of weeds varies so that not all weeds are contacted. If this occurs, repeat treatment up to the labeled rate may be necessary.

Shielded and Hooded Applicators: A shielded or hooded applicator directs the herbicide solution onto weeds while shielding desirable vegetation from the herbicide. Use nozzles that provide uniform coverage within the treated area. Keep shields on these sprayers adjusted to protect desirable vegetation. Exercise extreme care to avoid contact of the herbicide with desirable vegetation.

Wiper Applicators: Wiper applicators are devices that physically wipe appropriate amounts of this product directly onto the weed. Equipment must be designed, maintained and operated to prevent the herbicide solution from contacting desirable vegetation.

Adjust wiper applicators used over the top of desirable vegetation so that the wiper contact point is at least 2 inches above the desirable vegetation. Better results are obtained when more of the weed is exposed to the herbicide solution. Weeds should be a minimum of 6 inches above the desirable vegetation. Adjust the applicator height to ensure adequate contact with weeds as weeds not contacted by the herbicide solution will not be affected. Poor contact may occur when weeds are growing in dense clumps, in severe weed infestations, or when weed height varies dramatically. If this occurs, repeat treatment up to the labeled rate may be necessary.

Operate this equipment at ground speeds no more than 5 mph. Performance may be improved by reducing speed in areas of heavy weed infestations to ensure adequate wiper saturation. Better results may be obtained if two applications are made in opposite directions.

Droplets, mist, foam, or splatter of the herbicide settling onto desirable vegetation may result in discoloration, stunting or destruction. Avoid leakage or dripping onto desirable vegetation. Adjust height of applicator to ensure adequate contact with weeds. Keep wiping surfaces clean. Be aware that on sloping ground the herbicide solution may migrate, causing dripping on the lower end and drying of the wicks on the upper end of a wiper applicator.

Do not use wiper equipment when weeds are wet.

Mix only the amount of solution to be used during a one-day period as reduced activity may result from use of leftover solutions. Clean wiper parts by thoroughly flushing with water immediately after using this product.

For best results, use a nonionic surfactant at a rate of 10 percent by volume of total herbicide solution for all wiper applications.

Rope or Sponge Wick Applicators: Use solutions of 33 to 75 percent of this product in water.

Panel Applicator: Use solutions of 33 to 100 percent of this product in water.

Injection Systems

Equipment: Aerial or ground injection sprayers.

This product may be used in aerial or ground injection spray systems. It may be used as a liquid concentrate or diluted prior to injecting into the spray stream. Do not mix this product with the concentrate of other products when using injection systems.

Controlled Droplet Applicator (CDA)

Equipment: Hand-held or boom-mounted applicators that produce a spray consisting of a narrow range of droplet sizes.

The rate of this product applied per acre by vehicle-mounted CDA equipment must not be less than the amount specified on this label when applied by conventional broadcast equipment. For vehicle-mounted CDA equipment, apply 3 to 15 gallons of water per acre.

For the control of annual weeds with hand-held CDA units, apply a 20 percent solution of this product at a flow rate of 2 fl oz per minute and a walking speed of 1.5 mph (1 1/2 pints of product per acre). For control of perennial weeds, apply a 20 to 40 percent solution of this product at a flow rate of 2 fl oz per minute and a walking speed of 0.75 mph (3 to 6 pints of product per acre).

CDA equipment produces a spray pattern that is not easily visible. Exercise extreme care to avoid spray or drift contacting the foliage or any other green tissue of desirable vegetation as damage or destruction may result.

Use Sites

Use this product in noncrop areas, including airports, apartment complexes, aquatic sites, Christmas tree farms, commercial sites, Conservation Reserve Program (CRP) areas, ditch banks, driveways, dry ditches, dry canals, fencerows, golf courses, greenhouses, habitat management, industrial areas, lumber yards, manufacturing sites, municipal sites, natural areas, office complexes, ornamentals, parking areas, parks, pastures, petroleum tank farms and pumping installations, plant nurseries, public areas, railroads, rangeland, recreation areas, utility rights-of-way, roadsides, shadehouses, sod or turf seed farms, sports complexes, storage areas, substations, turfgrass areas, utility sites, warehouse areas, wildlife habitat management areas, and in grazed areas on these sites.

Limitations, Restrictions, and Exceptions

TURFGRASS RENOVATION, SEED OR SOD PRODUCTION

This product controls most existing vegetation prior to renovating turfgrass areas or establishing turfgrass grown for seed or sod. For maximum control of existing vegetation, delay planting or sodding to determine if any regrowth from escaped underground plant parts occurs. When repeat treatments are necessary, sufficient regrowth must be attained prior to application. For warm season turfgrass, including bermudagrass, summer or fall applications provide the best control. Where existing vegetation is growing under mowed turfgrass management, apply this product after omitting at last one regular mowing to allow sufficient growth for good interception of the spray.

Do not disturb soil or underground plant parts before treatment. Delay tillage or renovation techniques, including vertical mowing, coring, or slicing, for seven days after application to allow translocation into underground plant parts.

Desirable turfgrass may be planted following the above procedures.

Hand-held equipment may be used for spot treatment of unwanted vegetation growing in existing turfgrass. Broadcast or hand-held equipment may be used to control sod remnants or other unwanted vegetation after sod is harvested.

Do not feed or graze turfgrass grown for seed or sod production for eight weeks following application.

Perennial Weeds

Best results are obtained when perennial weeds are treated after they reach the reproductive stage of growth (seedhead initiation in grasses and bud formation in broadleaves). Best results are obtained when non-flowering plants are treated when they reach a mature stage of growth. In many situations, applications are required prior to these growth stages. Under these conditions, use a higher rate in the rate range.

When using spray to wet treatments with hand-held equipment, ensure thorough coverage of the plant. For best results, use a 1.5 percent solution on harder to control perennials including bermudagrass, dock, field bindweed, hemp dogbane, milkweed and Canada thistle.

Use a 4 to 7 percent solution of this product in low volume directed spray applications. Spray coverage should be uniform with at least 50 percent of the foliage contacted. For best results, cover the top one-half of the plant. To ensure adequate spray coverage, spray both sides of large or tall weeds when foliage is thick and dense or where there are multiple sprouts.

Allow 7 days or more after application before tillage.

Method

[Broadcast/Foliar Air](#)

[Broadcast/Foliar Ground](#)

[Directed Spray](#)

Rates

[field_rates 0](#)

[field_rates 1](#)

-

Restricted Entry Interval

4 hours

Timings

[Postemergence \(Weed\)](#)