

PERENNIAL GRASS SEED CROPS DURING THE YEAR OF ESTABLISHMENT - WASHINGTON - BILLBUGS AND CUTWORMS

General Information

Product Information

Lorsban Advanced insecticide is an emulsion in water for use in listed crops. This product resists washoff once it is dry. Target pests and application rates are provided in the accompanying tables.

Use Precautions and Restrictions

Insect control may be reduced at low spray volumes under high temperature and wind conditions.

Some reduction in insect control may occur under unusually cool conditions.

Flood irrigation: To avoid contamination of irrigation tail waters, do not flood irrigate within 24 hours following a soil surface or foliar application of Lorsban Advanced.

Do not aerially apply this product in Mississippi.

Insecticide Resistance Management (IRM)

Lorsban Advanced contains a Group 1B insecticide. Insect/mite biotypes with acquired resistance to Group 1B may eventually dominate the insect/mite population if Group 1B insecticides are used repeatedly in the same field or in successive years as the primary method of control for targeted species. This may result in partial or total loss of control of those species by Lorsban Advanced or other Group 1B insecticides.

To delay development of insecticide resistance:

- Avoid consecutive use of insecticides with the same mode of action (same insecticide group) on the same insect species.
- Use tank mixtures or premix products containing insecticides with different modes of action (different insecticide groups) provided the products are registered for the intended use.
- Base insecticide use upon comprehensive Integrated Pest Management (IPM)

programs.

- Monitor treated insect populations in the field for loss of effectiveness.
- Contact your local extension specialist, certified crop advisor, and or manufacturer for insecticide resistance management and/or IPM recommendations for the specific site and resistant pest problems.
- For further information or to report suspected resistance, you may contact Dow AgroSciences by calling 800-258-3033.

Application Directions

Broadcast Foliar Application

Apply with conventional power-operated spray equipment using nozzles and spray pressures specified for insecticides. Apply Lorsban Advanced in a spray volume of not less than 2 gallons per acre (gpa) for aerial application equipment (fixed wing or helicopter) or not less than 10 gpa for ground equipment, unless otherwise specified. Increase spray volume to ensure adequate coverage with increased density and height of crop canopy.

Ground Application: Orient the boom and nozzles so that uniform coverage is obtained. The swath width should not be wider than the boom. Follow nozzle manufacturer's specifications for insecticide nozzles with respect to nozzle type, pressure, and spacing.

Broadcast Soil Application

Apply with conventional power-operated spray equipment that will apply the product uniformly to the soil surface. Use nozzles that produce medium or coarse droplets (255 to 400 microns). Unless otherwise indicated, a spray volume of 10 gpa or more is needed. For band application, use proportionally less spray volume.

Aerial Application

Use a minimum spray volume of 2 gpa. Mark swaths by mechanical flagging, permanent markers or GPS equipment.

Chemigation Application

Apply Lorsban Advanced through properly equipped chemigation systems for insect control in alfalfa, almond (orchard floors only), citrus (orchard floors only), corn (field and sweet), cotton, cranberry, peppermint, sorghum, soybeans, spearmint,

sugarbeet, orchard floors (pecan and walnut), and wheat, or other crops as specified in Dow AgroSciences supplemental labeling. Do not apply this product by chemigation unless specified in crop-specific directions in this label or Dow AgroSciences supplemental labeling. Do not apply to labeled crops through any other type of irrigation system.

Note: Unless otherwise indicated in specific use directions, the application rates for chemigation are the same as those specified for broadcast application.

Limitations, Restrictions, and Exceptions

Apply Lorsban Advanced to perennial grass seed crops, such as Kentucky bluegrass, orchardgrass, tall fescue, wheat grass, timothy grass, and perennial ryegrass, to control billbugs, cutworms, and aphids during the year of establishment.

Billbugs and Cutworms: Apply 1 quart of Lorsban Advanced per acre in 20 gallons of water per acre as a broadcast spray using power-operated ground spray equipment. For aerial application, apply in 2 to 5 gallons of water per acre. For improved results, thatch should be moist at time of application and the insecticide should be washed into the thatch immediately after application. If sprinkler irrigation is available, apply 1/4 to 1/2 inch of water after application to help incorporate the material, or apply to early morning dew-covered fields, or during or immediately following a light rain. For billbug control, apply in early April when adult billbugs are active. Control of billbugs is dependent upon proper application timing and coverage. For cutworm control, apply in either spring or fall when larvae or larval damage first appears. Fall applications should follow sufficient irrigation and/or rainfall.

Specific Use Restrictions:

- Restricted Entry Interval: Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 24 hours unless PPE required for early entry is worn.
- Maximum single application rate is 0.94 lb ai chlorpyrifos (2 pints of Lorsban Advanced) per acre. Do not make more than two applications of chlorpyrifos per year when using 0.94 lb ai chlorpyrifos (2 pints of Lorsban Advanced) per acre.
- Do not apply more than 2.5 lb ai chlorpyrifos (5 1/3 pints of Lorsban Advanced) per acre per year. Do not make more than three applications of chlorpyrifos per year when using 1 3/4 pints or less of Lorsban Advanced per treatment.
- Do not re-treat with Lorsban Advanced or other product containing chlorpyrifos

within 10 days of the previous application.

- Under this SLN label, use of Lorsban Advanced is deemed a non-food use by the Washington Department of Agriculture and, as such, is not for use on fields producing grass for livestock feed.
- Chemigation: For use under this SLN label, do not apply this product through any type of irrigation system.
- Lorsban Advanced may only be applied to grass seed fields during the year of establishment.
- Do not apply this insecticide within one year (365 days) of harvesting grass seed.
- Do not graze livestock or harvest forage from treated fields within one year (365 days) of application.
- Pollinator Protection: This product is highly toxic to bees exposed to direct treatment or residues on blooming crops or weeds. Do not apply this product to blooming grass seed during the pollination period. Apply as a pre-bloom spray, or apply at the end of the pollination period. Bee colonies used for pollination should be removed from the field being treated prior to the application. Do not apply this product or allow it to drift to blooming crops or weeds if bees are visiting the treatment area. Notify beekeepers pollinating crops within 1/4 mile of the field to be treated at least 48 hours prior to the application. Consult your cooperative agricultural extension service for additional information concerning billbug, cutworm, and aphid control practices in your area.

Method

[Broadcast/Foliar Air](#)

[Broadcast/Foliar Ground](#)

[Broadcast/Foliar Air](#)

[Broadcast/Foliar Ground](#)

Rates

[field_rates 0](#)

[field_rates 1](#)

•

Restricted Entry Interval

24 hours

Timings

[Billbug: In early April when adult billbugs are active.](#)

[Billbug: In early April when adult billbugs are active.](#)