

# **SUBTERRANEAN TERMITES - CARTON AND NESTS**

## General Information

### APPLICATION INSTRUCTIONS

This product may be used alone to kill subterranean or drywood termites or as a supplement to a post-construction treatment of Termidor 80 WG Termiticide/Insecticide, Termidor HE High-Efficiency Termiticide, or Termidor SC Termiticide/Insecticide to provide protection from subterranean termites. This product will kill workers and/or winged reproductive forms of subterranean and/or drywood termites present at the time of application and will provide residual control of termites contacting the treated surfaces only as described in the Directions For Use. This product is intended only as a spot treatment.

Application of this product will not provide structural protection nor substitute for mechanical alteration or soil or foundation treatment with registered termiticide products. This product is not to be used as sole protection against termites, as it is not intended for elimination of termite infestations nor for protection against future infestations.

### APPLICATION DEVICE

This ready-to-use product is contained in a nonrefillable package and is for use with a specially designed applicator. Approximately 3 bulb compressions of the applicator delivers 0.1 gram of product.

### APPLICATION SITES

Apply into voids and galleries, shelter tubes, termite nests or other inaccessible areas of buildings, trees, utility poles, fencing, decking materials, railroad trestles, piers, beams and other structural or landscape timbers where termite damage is observed or termite activity is present or suspected.

If necessary, gain access to a gallery or void by creating an opening into it. Openings should be approximately every 24 inches or into each void.

Multiple injection points for each void, carton, or gallery may be required. DO NOT

apply more than 5 grams product (25.0 milligrams fipronil a.i.) per 1,000 square feet of total internal surface area of the void, carton, or gallery.

## REAPPLICATION INTERVAL

This product may be applied to previously treated sites if termite activity is found 30 days after previous application.

## Limitations, Restrictions, and Exceptions

### Subterranean Termites:

For applications into cartons and nests, apply 0.3 to 3 grams per injection point.

### Method

[Injection](#)

### Rates

[field\\_rates 0](#)

•

### Timings

[N.A.](#)