

## **TREES AND SHRUBS (OAK)**

### General Information

#### **EIGHT INSECT CONTROL VEGETABLE, FRUIT & FLOWER CONCENTRATE**

Do not water the treated area to the point of run-off. Do not make applications during rain. All outdoor applications must be limited to spot or crack-and-crevice treatments only, except for the following permitted uses:

(1) Treatment to soil or vegetation around structures; (2) Applications to lawns, turf, and other vegetation; (3) Applications to building foundations, up to a maximum height of 3 feet.

Other than applications to building foundations, all outdoor applications to impervious surfaces such as sidewalks, driveways, patios, porches and structural surfaces (such as windows, doors, and eaves) are limited to spot and crack-and-crevice applications only. Application is prohibited directly into sewer drains, or to any area like a gutter where drainage to sewers, storm drains, water bodies, or aquatic habitat can occur. Do not allow the product to enter any drain during or after application.

For lawns and vegetables, measure the area to be sprayed. Calculate square feet by multiplying length by width. For all other applications, apply as a thorough cover spray. Use a clean sprayer. Carefully measure and mix the amount of concentrate and water as directed. Spray as directed. Thorough coverage is important. Flush sprayer with clean water after each use. Make applications when insects first appear. Repeat applications as directed. See directions for complete insect list. People and pets should not be allowed in treated areas until spray has dried.

Use dilution rates as indicated.

1 fl. oz. = 6 tsp. or 2 Tbsp.

8 fl. oz. = 1/2 pt. or 1 cup

### Limitations, Restrictions, and Exceptions

#### **TREES & SHRUBS**

How to use:

Use 1 fl. oz. in 1 gal. of water. Apply as a thorough spray, wetting leaves and branches to the dripping point. Try to penetrate dense foliage. Spray in the late afternoon or evening, when the temperature ranges from 50° to 75°F and when there is little or no wind. Spray at the first sign of insects. Repeat as necessary; use intervals of 4-8 days. Application can be made up to the day of collection.

Method

[Spray](#)

Rates

[field\\_rates 0](#)

•

Timings

[When the insects first appear.](#)

[When the insects first appear.](#)