

## **FRUIT AND NUT TREES (APPLES)**

### General Information

#### EIGHT INSECT CONTROL VEGETABLE, FRUIT & FLOWER CONCENTRATE

Do not water the treated area to the point of run-off. Do not make applications during rain. All outdoor applications must be limited to spot or crack-and-crevice treatments only, except for the following permitted uses:

(1) Treatment to soil or vegetation around structures; (2) Applications to lawns, turf, and other vegetation; (3) Applications to building foundations, up to a maximum height of 3 feet.

Other than applications to building foundations, all outdoor applications to impervious surfaces such as sidewalks, driveways, patios, porches and structural surfaces (such as windows, doors, and eaves) are limited to spot and crack-and-crevice applications only. Application is prohibited directly into sewer drains, or to any area like a gutter where drainage to sewers, storm drains, water bodies, or aquatic habitat can occur. Do not allow the product to enter any drain during or after application.

For lawns and vegetables, measure the area to be sprayed. Calculate square feet by multiplying length by width. For all other applications, apply as a thorough cover spray. Use a clean sprayer. Carefully measure and mix the amount of concentrate and water as directed. Spray as directed. Thorough coverage is important. Flush sprayer with clean water after each use. Make applications when insects first appear. Repeat applications as directed. See directions for complete insect list. People and pets should not be allowed in treated areas until spray has dried.

Use dilution rates as indicated.

1 fl. oz. = 6 tsp. or 2 Tbsp.

8 fl. oz. = 1/2 pt. or 1 cup

### Limitations, Restrictions, and Exceptions

#### FRUIT & NUT TREES

## GENERAL DIRECTIONS

Use 2 fl. oz. in 1 gal. of water. Apply when insects appear. Wet the plants to the dripping point. Try to hit underside of leaves and penetrate dense foliage.

How to use:

Repeat as required to maintain control. Do not make more than 3 applications. Do not apply after petal fall.

Method

[Spray](#)

Rates

[field\\_rates 0](#)

•

Timings

[When insects appear.](#)