

FRUIT & NUT CROPS - PEACHES - CORYNEUM BLIGHT (PEACH BLIGHT, SHOTHOLE)

General Information

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

FOR OUTDOOR RESIDENTIAL USE ONLY.

Do not allow this product to drift. Do not apply this product in a way that will contact any person or pet, either directly or through drift. Keep people and pets out of the area during application.

ENTRY RESTRICTIONS

Do not allow people or pets to enter the treated area until sprays have dried.

COMPATIBILITY

This product is compatible with most commonly used insecticides and fungicides, except highly alkaline materials such as hydrated lime. Captan should not be used in combination with or closely following oil sprays. Observe all cautions and limitations on labeling of all products used in mixture.

Where foliage applications are directed, mix designated amount of material in a little water to make a thin paste, add to specified amount of water, and keep mixture agitated during application. Use a compression type or hose-end sprayer.

Limitations, Restrictions, and Exceptions

FRUIT & NUT CROPS

Size:

Young or Dwarf: Amount of Spray Needed Per Tree Diluted Spray: 1 to pints

Medium: Amount of Spray Needed Per Tree Diluted Spray: 1 to 2 quarts

Large: Amount of Spray Needed Per Tree Diluted Spray: ½ to 3 gallons

PEACHES

Coryneum blight (peach blight, shothole) – Apply in pink, bud, full bloom, petal fall, and cover sprays as necessary, and as a post harvest spray (before leaves drop). Preharvest spray may be applied up to day of harvest.

Method

[Spray](#)

Rates

[field_rates 0](#)

•

Timings

[In pink, bud, full bloom, petal fall, and cover sprays as necessary, and as a post harvest spray \(before leaves drop\).](#)