

## **FRUIT & NUT CROPS - GRAPES (U.S. EXCEPT CA)**

### General Information

### DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

### FOR OUTDOOR RESIDENTIAL USE ONLY.

Do not allow this product to drift. Do not apply this product in a way that will contact any person or pet, either directly or through drift. Keep people and pets out of the area during application.

### ENTRY RESTRICTIONS

Do not allow people or pets to enter the treated area until sprays have dried.

### COMPATIBILITY

This product is compatible with most commonly used insecticides and fungicides, except highly alkaline materials such as hydrated lime. Captan should not be used in combination with or closely following oil sprays. Observe all cautions and limitations on labeling of all products used in mixture.

Where foliage applications are directed, mix designated amount of material in a little water to make a thin paste, add to specified amount of water, and keep mixture agitated during application. Use a compression type or hose-end sprayer.

### Limitations, Restrictions, and Exceptions

### FRUIT & NUT CROPS

#### Size:

Young or Dwarf: Amount of Spray Needed Per Tree Diluted Spray: 1 to pints

Medium: Amount of Spray Needed Per Tree Diluted Spray: 1 to 2 quarts

Large: Amount of Spray Needed Per Tree Diluted Spray: ½ to 3 gallons

## GRAPES (U.S. except CA)

Phomopsis cane and leaf spot, downy mildew, suppression of black rot – Apply sufficient spray volume to provide thorough, even coverage when shoots are ½ to 1½ inches long, when shoots are 3 to 5 inches long and when shoots are 9 to 12 inches long. Repeat just before bloom, immediately after bloom, and continue at 10 to 14 day intervals as needed. Use the lower rate when spraying less susceptible grape varieties or when conditions are less favorable for disease development. Use the higher rate on susceptible grape varieties and during periods of weather highly favorable for disease development. May be applied up to harvest.

### Method

[Spray](#)

Rates

[field\\_rates 0](#)

- 

### Timings

[When shoots are ½ to 1½ inches long, when shoots are 3 to 5 inches long and when shoots are 9 to 12 inches long.](#)