

ORNAMENTALS - CONTROL OF FUNGAL DISEASES

General Information

- For use on houseplants, roses, flowers, ornamental trees and shrubs, fruits, nuts and vegetables
- Controls insects, fungal diseases and mites, including spider mites

BEFORE YOU USE

Use Precautions:

- Some varieties of grapes and cucurbits are sensitive to sulfur and some plant damage may occur.
- Use care or do not spray on sulfur sensitive plants. If not sure, test spray a few leaves 2 days before spraying the entire plant.

Use Restrictions:

- Do not apply to strawberry fruit that will be used for canning.
- Do not wet plants to point of runoff or drip.
- Do not water the treated areas to point of runoff or drip.
- Do not make applications during rain.
- Not for use in outdoor residential misting systems.
- Do not apply more than 10 times per season.
- Do not reapply within 3 days except under extreme pest pressure.
- In case of extreme pest pressure do not reapply within 24 hours.
- Do not apply more than 1 time per day.
- Do not apply more than 18.5 fl. oz. per 100 square feet per application.
- Do not use if temperatures are expected to be over 90°F.
- Do not apply this product in a way that will contact adults, children or pets, either directly or through drift.
- Remove pets, birds, and cover aquariums before spraying.
- Remove or cover exposed food and drinking water before application. Remove or cover dishes, utensils, food processing equipment, and food preparation surfaces, or wash them before use.
- Do not apply directly to or near water, storm drains, or drainage ditches. Do not apply when windy. To prevent product runoff, do not over water treated area or

apply prior to heavy rainfall.

- Do not apply directly into sewers or drains, or to any area like a gutter where drainage to storm sewers, water bodies, or aquatic habitat can occur. Do not allow the product to enter any drain during or after application.
- Do not allow adults, children or pets to enter the treated area until sprays have dried.
- Do not spray plants with this product within 21 days of an oil spray.

Limitations, Restrictions, and Exceptions

FOR CONTROL OF FUNGAL DISEASES

Thoroughly spray all areas of the plant, especially new shoots. For best control apply as a protective spray early in the season before the diseases are noticed.

Re-spray every 7 to 10 days or after rain. May be used up to day before harvest.

Method

[Spray](#)

Timings

[Early in the season before the diseases are noticed.](#)