

SOYBEANS - NOT LESS THAN 15 DAYS PRIOR TO PLANTING

General Information

USE RESTRICTIONS

Do not apply this product through any type of irrigation system.

Do not use in or near a greenhouse.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application.

Do not apply to frozen or snow covered soil.

Do not apply to farm alleys or roads where traffic may result in treating dust settling onto crops or other desired vegetation.

Do not apply within 300 yards of non-dormant pears.

Post directed and layby application of this product should only be applied to healthy growing crops.

Do not apply to powdery soils that are susceptible to wind displacement unless irrigation can be applied immediately after application.

Read and follow all directions, restrictions and precautions on this label and on the labels of any products for which a tank mixture is being considered.

PRODUCT INFORMATION

INJURY TO CROPS FROM THIS HERBICIDE MAY OCCUR. IF YOU ARE NOT PREPARED TO ACCEPT SOME DEGREE OF CROP INJURY Do not USE THIS PRODUCT.

Crop varieties vary in response to 2,4-D and some are easily injured. Apply this product only to varieties known to be tolerant to 2,4-D. If you are uncertain concerning tolerant varieties or local use situations that may affect crop tolerance to 2,4-D, consult your seed company, State Agricultural Extension Service or qualified crop consultant for advice.

Be sure that use of this product conforms to all applicable laws, rules and regulations. Certain states have restrictions pertaining to application distances from susceptible crops. The applicator should become familiar with these laws, rules or regulations and follow them exactly.

APPLICATION PROCEDURES

Apply by air or ground equipment in sufficient gallonage to obtain adequate coverage, except as otherwise directed on this label. Use 2 or more gallons of water per acre for aerial application and 10 or more gallons of water per acre for ground application.

RESISTANCE MANAGEMENT

This product contains active ingredients in Groups 4 and 14. Any weed population may contain or develop plants that are resistant to this product and other Group 4 and 14 herbicides. Weed species with acquired resistance to Group 4 and 14 herbicides may eventually dominate the weed population if Group 4 and 14 herbicides are used repeatedly in the same application area or in successive years as the primary method of control for targeted species. This may result in partial or total loss of control of those species by this product or other Group 4 and 14 herbicides.

To delay or prevent herbicide resistance consider the following practices:

Avoid the use of more than two consecutive applications of this product or other herbicides that have a similar target site of action.

Alternate herbicides used for weed control.

Base herbicide use on a comprehensive Integrated Pest Management (IPM) program.

Monitor treated weed populations for loss of efficacy.

Contact your local extension specialist, other experts appropriate to aquatic use, and/or manufacturer for resistance and/or integrated weed management practices.

For further information or to report suspected resistance, you may contact Innvictis.

Limitations, Restrictions, and Exceptions

SOYBEANS

(Preplant Only)

DIRECTIONS

Apply not less than 15 days prior to planting soybeans, when weeds are actively growing.

In addition to those weeds found on the WEED LIST, this product will suppress or control the following broadleaf weeds frequently encountered in reduced tillage soybean production systems: alfalfa*, bullnettle, smallflowered bittercress, Carolina geranium, smallflowered buttercup, common and rough cinquefoil, red clover*, horseweed or maretail, mousetail, wild mustard, field pennycress, cutleaf evening

primrose, common purslane, speedwell, velvetleaf, and Virginia copperleaf. * These weeds are only partially controlled.

Apply no more than 2.0 pints of this product in one season prior to planting soybeans. After applying, plant soybean seed as deep as practical or at least 1-1/2 to 2 inches deep. Adjust the planter press wheel, if necessary, to ensure that planted seed is completely covered.

Applications of 2 pints per acre will provide residual control as described in the Table - Broadleaf Weeds Controlled by Residual Activity of This Product.

If desired, this product may be applied pre-plant to soybeans in tank mixtures with other herbicides such as Poast, Poast Plus, Roundup, Roundup D-Pak, Honcho, Gramoxone Extra, Prowl, Pursuit Plus, Scepter, Scepter 70 DG, Squadron and others that are registered for pre-plant soybean use.

Do not tank mix this product with acetochlor (Warrant), alachlor (Micro-Tech), flufenacet (Axiom, Domain), metolachlor (Dual Magnum, Dual II Magnum, Boundary) or dimethenamid (Frontier or Outlook) within 15 days of planting soybeans, unless soybeans are planted under no-till or minimum tillage conditions on wheat stubble or no-till field corn stubble.

NOTE: Unacceptable injury to soybeans planted in fields previously treated with this product may occur and the extent of injury will depend on weather and agronomic factors such as the amount of weed vegetation and previous crop residue present that may be in effect between the time of application and the emergence of the soybean plant.

RESTRICTIONS AND LIMITATIONS FOR USE IN SOYBEANS (PREPLANT)

- Do not apply this product when weather conditions such as temperature, air inversions, or wind favor drift from treated areas to susceptible plants.
- Apply no more than 2.0 pints (1.0 lb. 2,4-D ae) of this product per acre in one season prior to planting soybeans.
- Only one application per growing season, regardless of the application rate used, is allowed.
- Do not apply this product prior to planting soybeans if you are not prepared to accept the results of soybean injury including possible loss of stand and yield.
- Do not replant fields treated with this product in the same growing season with crops other than those labeled for 2,4-D pre-plant use.
- Do not mow or cultivate weeds prior to treating with this product as poor control may result.
- Do not apply this product pre-plant to soybeans in fields having a coarse-textured soil where the percent organic matter is <1.0%.

- Do not irrigate when soybeans are cracking.
- Do not graze treated fields or feed treated hay to livestock.

Method

[Broadcast/Foliar Air](#)

[Broadcast/Foliar Ground](#)

[Broadcast/Foliar Air](#)

[Broadcast/Foliar Ground](#)

Rates

[field_rates 0](#)

-

Restricted Entry Interval

48 hours

Timings

[Preplant](#)

[Postemergence \(Weed\)](#)