

HOPS

General Information

USE RESTRICTIONS

Do not apply this product through any type of irrigation system.

Do not use in or near a greenhouse.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application.

Do not apply to frozen or snow covered soil.

Do not apply to farm alleys or roads where traffic may result in treating dust settling onto crops or other desired vegetation.

Do not apply within 300 yards of non-dormant pears.

Post directed and layby application of this product should only be applied to healthy growing crops.

Do not apply to powdery soils that are susceptible to wind displacement unless irrigation can be applied immediately after application.

Read and follow all directions, restrictions and precautions on this label and on the labels of any products for which a tank mixture is being considered.

PRODUCT INFORMATION

INJURY TO CROPS FROM THIS HERBICIDE MAY OCCUR. IF YOU ARE NOT PREPARED TO ACCEPT SOME DEGREE OF CROP INJURY Do not USE THIS PRODUCT.

Crop varieties vary in response to 2,4-D and some are easily injured. Apply this product only to varieties known to be tolerant to 2,4-D. If you are uncertain concerning tolerant varieties or local use situations that may affect crop tolerance to 2,4-D, consult your seed company, State Agricultural Extension Service or qualified crop consultant for advice.

Be sure that use of this product conforms to all applicable laws, rules and regulations. Certain states have restrictions pertaining to application distances from susceptible crops. The applicator should become familiar with these laws, rules or regulations and follow them exactly.

APPLICATION PROCEDURES

Apply by air or ground equipment in sufficient gallonage to obtain adequate coverage, except as otherwise directed on this label. Use 2 or more gallons of water per acre for aerial application and 10 or more gallons of water per acre for ground application.

RESISTANCE MANAGEMENT

This product contains active ingredients in Groups 4 and 14. Any weed population may contain or develop plants that are resistant to this product and other Group 4 and 14 herbicides. Weed species with acquired resistance to Group 4 and 14 herbicides may eventually dominate the weed population if Group 4 and 14 herbicides are used repeatedly in the same application area or in successive years as the primary method of control for targeted species. This may result in partial or total loss of control of those species by this product or other Group 4 and 14 herbicides.

To delay or prevent herbicide resistance consider the following practices:

Avoid the use of more than two consecutive applications of this product or other herbicides that have a similar target site of action.

Alternate herbicides used for weed control.

Base herbicide use on a comprehensive Integrated Pest Management (IPM) program.

Monitor treated weed populations for loss of efficacy.

Contact your local extension specialist, other experts appropriate to aquatic use, and/or manufacturer for resistance and/or integrated weed management practices. For further information or to report suspected resistance, you may contact Innvictis.

Limitations, Restrictions, and Exceptions

HOPS

DIRECTIONS

Make directed applications to the row middles. Make up to 3 applications at 30-day intervals with the last application before harvest.

RESTRICTIONS AND LIMITATIONS FOR HOPS

- Limited to 3 applications per crop cycle.
- Maximum of 1 pint (0.5 lbs. 2,4-D ae) per acre per application.
- Maximum of 3 pints (1.5 lbs. 2,4-D ae) per acre per crop cycle.
- Minimum of 30 days between applications.
- (PHI) Do not harvest within 30 days of application.

- Do not use with an adjuvant.
- Do not allow spray to contact green stem, foliage, flowers or cones or unacceptable injury may occur.
- Do not mow or rake over treated areas, as dust created by mowing may drift onto sensitive crops or vegetation resulting in injury.

Method

[Broadcast/Foliar Air](#)

[Broadcast/Foliar Ground](#)

Pre-Harvest Interval

30 days

Rates

[field_rates 0](#)

•

Restricted Entry Interval

48 hours

Timings

[N.A.](#)