

BERRY AND SMALL FRUITS (EPA GROUPING 11) - SOIL APPLIED

General Information

GENERAL RECOMMENDATIONS

Before including any other fertilizer with LEVITATE in seed furrow applications, conduct a jar test of the products to be mixed in their respective proportions and observe it for 24 hours before mixing large quantities. The person making the recommendation or application is responsible for verifying the seed safety of any products mixed with LEVITATE. When used as directed this product does not supply all nutrients required by plants and is to supplement a soil fertility program based on soil tests.

SOIL OR FOLIAR APPLICATIONS: Consult rate chart below. Water may be added to dilute LEVITATE when higher volume application is desired. Optimum rate of application will vary between fields, depending on soil pH and organic matter content. For best results, follow soil test or plant analysis recommendations.

CROP SAFETY: This product has been tested on a number of crop varieties with no phytotoxicity observed at labeled rates; however, this product has not been tested on all varieties and cultivars. Due to the large number of plant species and their associated varieties or cultivars, and due to variable growing conditions, it is impossible to test every plant for tolerance to this product. It is the responsibility of the user to determine if this product can be used safely prior to commercial use. In a small test area, test the specified rates on plants for phytotoxicity prior to general use. Before using any tank mix, test the combination on a small portion of the crop to be treated to ensure that a phytotoxic response will not occur as a result of application. Treat a small group of target crop including plant cultivars and evaluate for 3-7 days prior to making an application to your entire crop.

Limitations, Restrictions, and Exceptions

1-2.5 gal/acre per application as needed

Method

[Soil application](#)

Rates

[field_rates 0](#)

-

Timings

[N.A.](#)