

GOLF COURSE TURFGRASS: TURF-PARASITIC NEMATODES - 14 TO 21 DAY INTERVAL

General Information

PRODUCT INFORMATION

- Divanem is a suspension concentrate containing abamectin which, when applied according to directions for use, will control turf-parasitic nematodes such as sting (*Belonolaimus longicaudatus*), lance (*Hoplolaimus galeatus*), root-knot (*Meloidogyne* spp.), and ring (*Criconemoides* spp.) on golf courses and professional and collegiate sports fields.
- Divanem should be applied as a seasonal program.
- Apply Divanem as an early curative treatment (after appropriate nematode extraction, identification, and counts).
- Multiple applications may be required before improvements in turf quality are observed.
- Maximum annual rate must not exceed 0.27 lb abamectin/calendar year or 50 fl oz Divanem/A/calendar year.

APPLICATION DIRECTIONS

Application Equipment

For best results, use spray nozzles that produce extremely coarse droplets [such as Turbo Floodjet (TF5) or TurfJet (TTJ10)].

RESTRICTIONS AND PRECAUTIONS

Use Restrictions

- DO NOT apply to golf course roughs.
- DO NOT apply to residential turf, sports fields (other than professional or collegiate), or commercial turf.
- DO NOT use aerial application.
- Chemigation: DO NOT apply this product through any type of irrigation system.
- DO NOT apply through an ultra-low volume spray system.
- DO NOT apply to turf under heat or moisture stress.
- DO NOT GRAZE TREATED TURF OR FEED TURF CLIPPINGS FROM ANY TREATED AREA TO POULTRY OR LIVESTOCK.

Limitations, Restrictions, and Exceptions

GOLF COURSE TURFGRASS

Turfgrass (including all cultivars, varieties, and/or hybrids)

Application Timing

- Apply at this rate on a 14- to 21-day interval.

Use Directions

- Apply in the early morning while grass is wet with dew or irrigate the area prior to application with 0.1 inches of water. Spray onto wet turf.
- Irrigate with 0.1 to 0.5 inches of water beginning within 1 hour of application to move Divanem through the thatch. For best results, irrigate before the spray droplets have dried on the turf.
- Apply in 2 gallons of water per 1,000 square feet of turf. When lower application volumes are used, irrigating in should begin as soon as possible after application.
- Combination treatments of Divanem and a fungicide, such as Heritage Action, Heritage Fungicide, or Heritage TL, are recommended to reduce fungal infections following nematode feeding. Heritage Action, Heritage Fungicide, or Heritage TL should be watered in with Divanem.
- Divanem is also recommended in combination with a foliar fungicide program which includes Daconil Action. Daconil Action should be applied separately and not watered in.
- Directions For Use and Use Restrictions must be followed for all products being applied.

USE RESTRICTIONS

- 1) Maximum Single Application Rate: 0.067 lb abamectin/A (12.2 fl oz Divanem /A)
- 2) Minimum Application Interval: 14 days
- 3) Maximum Annual Rate: 0.27 lb abamectin/calendar year (50 fl oz Divanem/A/calendar year)

Method

[Spray](#)

Rates

[field_rates 0](#)

•

Timings

[In the early morning while grass is wet with dew](#)