

BETWEEN CROP APPLICATIONS - PERENNIAL (OTHER PERENNIALS)

Limitations, Restrictions, and Exceptions

BETWEEN CROP APPLICATIONS

PREPLANT DIRECTIONS (POSTHARVEST, FALLOW, CROP STUBBLE, SET-ASIDE) FOR BROADLEAF WEED CONTROL

VERITAS LV can be applied either postharvest in the fall, spring, or summer during the fallow period or to crop stubble/set-aside acres. Apply VERITAS LV as a broadcast or spot treatment to emerged and actively growing weeds after crop harvest (postharvest) and before a killing frost or in the fallow cropland or crop stubble the following spring or summer.

See Crop-Rotational Restrictions in section V. Restrictions and Limitations for the recommended interval between application and planting to prevent crop injury.

Rates and Timings:

Apply 4 - 32 fluid ounces of VERITAS LV per acre. Refer to Table 2 to determine use rates for specific targeted weed species. For best performance, apply VERITAS LV when annual weeds are less than 6" tall, when biennial weeds are in the rosette stage and to perennial weed regrowth in late summer or fall following a mowing or tillage treatment. The most effective control of upright perennial broadleaf weeds such as Canada thistle and Jerusalem artichoke occurs if VERITAS LV is applied when the majority of weeds have at least 4 - 6" of regrowth or for weeds such as field bindweed and hedge bindweed that are in or beyond the full bloom stage.

Avoid disturbing treated areas following application. Treatments may not kill weeds that develop from seed or underground plant parts such as rhizomes or bulblets, after the effective period for VERITAS LV. For seedling control, a follow-up program or other cultural practices could be instituted. For small grain in-crop uses of VERITAS LV, refer to the small grain section for details.

NOTES:

- DO NOT broadcast apply more than 32 fluid ounces per acre for single application.

Use the higher level of listed rate ranges when treating dense vegetative growth or perennial weeds with well-established root growth. "Other perennials" are defined as those listed in Table 1. General Weed List, Including ALS- and Triazine-Resistant Biotypes without footnote 1. The use on other perennials and on Woody Brush and Vine stems and for stem suppression is not registered in California.

- Rates higher than 32 fluid ounces per acre are for spot treatment only.

Method

[Broadcast/Foliar Air](#)

[Broadcast/Foliar Ground](#)

[Spot treatment](#)

[Broadcast/Foliar Air](#)

[Broadcast/Foliar Ground](#)

[Spot treatment](#)

[Broadcast/Foliar Air](#)

[Broadcast/Foliar Ground](#)

[Spot treatment](#)

Rates

[field rates 0](#)

•

Timings

[Post-harvest](#)

[Preplant](#)

[Postemergence \(Weed\)](#)