

# **ORANGES AND GRAPEFRUIT (BEARING) (FLORIDA AND TEXAS ONLY)**

## General Information

### USE RESTRICTIONS

Do not apply in spray solution above pH 10.

It is best to use COMITE II at pH 7 or lower.

Unless the rotated crop is a registered use for propargite, the following applies:

Do not plant any root crop in rotation within 6 months after last application of propargite to labeled crops.

Do not plant other crops in rotation within 2 months after last application of propargite to labeled crops.

Registered-labeled field and vegetable uses for propargite are:

Beans, dry Jojoba Seed crops only:

Corn (sweet, Mint Alfalfa field and popcorn) Peanuts Carrot, Cotton Potatoes Clover, Hops Sorghum Sugar Beets

### GENERAL INSTRUCTIONS

COMITE II is a liquid emulsifiable concentrate for preparing sprays to control mites. COMITE II is not systemic in action, therefore complete coverage of both upper and lower leaf surfaces and of fruit is necessary for effective control. COMITE II can be used in concentrate sprays as directed by label.

COMITE II performance is best when day temperatures average above 70°F.

### COMPATIBILITIES

Do not use penetrating surfactants. Under certain conditions a non-penetrating surfactant can be added. Rely on your prior use experience or consult your local surfactant supplier for specific recommendations.

Do not apply in combination with petroleum based foliar spray oils as phytotoxicity may occur.

COMITE II is not compatible with alkaline materials such as lime, lime sulfur or Bordeaux. The effectiveness of COMITE II will be reduced.

Except as noted, COMITE II is compatible with many pesticides.

Observe all cautions and limitation of labeling of all products used in mixtures.

However, due to variations in water quality, e.g. hardness and pH, it is required that users conduct small scale trials under local conditions to ensure compatibility prior

to large scale use.

Contact your supplier or the company representative for additional information on compatibility.

## Limitations, Restrictions, and Exceptions

### ORANGES & GRAPEFRUIT

Apply high rate to areas with high mite infestation and areas which historically have had high mite infestations.

Do not apply petroleum oils 30 days prior to or 40 days after applying COMITE II as phytotoxicity may occur on leaves and or fruit. Do not apply COMITE II mixed with petroleum oils or any material containing petroleum based solvents as phytotoxicity may occur on leaves and/or fruit.

Do not mix COMITE II with or apply prior to, or after, alkaline materials such as lime, Bordeaux mixture or lime sulfur.

Avoid COMITE II applications to immature leaves and fruit, and prevent spray solution from puddling, dripping, or running off leaves and/or fruit.

See COMPATIBILITIES section.

Ground Application: Apply in a minimum of 25 gallons total volume per acre. Apply in sufficient water for thorough coverage.

Aerial Application: Apply in a minimum of 10 gallons total volume per acre. Apply in sufficient water for thorough coverage.

ORANGE & GRAPEFRUIT RESTRICTIONS: Apply no more than 87 fl. oz. (4.1 lbs. propargite a.i.) per acre per year. Do not make more than 2 applications per season; the minimal spray interval is 28 days. See AGRICULTURAL USE REQUIREMENTS for general REI exemptions. There is no exception allowed for hand weeding.

Pre-harvest Interval: Do not harvest within 21 days of the last application.

### Method

[Broadcast/Foliar Air](#)

[Broadcast/Foliar Ground](#)

Pre-Harvest Interval

21 days

Rates

[field\\_rates 0](#)

- 

Restricted Entry Interval

20 days

Timings

[N.A.](#)