

INSECTICIDAL SOAP CONCENTRATE

General Information

INSECTICIDAL SOAP CONCENTRATE

- CONTROLS: ADELGIDS, APHIDS, EARWIGS, GRASSHOPPERS, PSYLLIDS, SAWFLY LARVAE, SCALE INSECTS, TENT CATERPILLARS, THRIPS, WHITEFLY, WOOLLY APHIDS AND OTHER LISTED PESTS.

- FOR USE ON VEGETABLES, FRUIT AND NUT TREES, CITRUS, BERRIES, ORNAMENTALS, SHRUBS, FLOWERS, AND TREES IN HOMES, GARDENS AND GREENHOUSES.

May be used on edible crops up to and including day of harvest after a minimum spray interval of 12 hours.

PESTS/SITES AND APPLICATION INSTRUCTIONS

This product is a contact spray and pests must be sprayed directly to achieve control. This product will control listed insect pests such as: adelgids (woolly aphids), aphids, lacebugs, mealybugs, mites, grasshoppers, leafhoppers, scale insects, plant bugs, sawfly larvae (pear and rose slugs), psyllids, tent caterpillars, thrips, spider mites, earwigs and whitefly.

Apply a 2% diluted product using hand-held trigger sprayers, hand sprayers or backpack sprayers. Thoroughly spray insect pests on plants to wetting with diluted product. Repeat treatment weekly or bi-weekly if needed. This product is a contact spray and is effective in its liquid state. Efforts should be made to thoroughly reach all plant parts including undersides of leaves.

For use indoors, outdoors and in greenhouses on fruit and nut trees (e.g., apples, cherries, peaches, pears, pecans, almonds); corn; soybeans; melons; tomatoes; vegetables (e.g., beans, cabbage, cucurbits, peas, potatoes); figs; small fruits (e.g., grapes, strawberry, raspberry); citrus; houseplants, flowering, foliage and bedding plants; ornamental and shade trees (e.g., birch, evergreens, holly, oak).

USE RESTRICTIONS

Do not use on sweet peas, nasturtiums, delicate ferns.

Do not spray when plants are under stress. Use with care on new seedlings, transplants and blooms. Do not spray during full sun. Spray early in morning or evening or when overcast. Soap spray may cause marking of some varieties.

If concerned about sensitivity of plants, apply to individual plants or small areas of plants and wait to determine if plant damage (phytotoxicity) occurs prior to treatment of larger areas.

Method

[Directed Spray](#)

Rates

[field_rates 0](#)

[field_rates 1](#)

[field_rates 2](#)

-

Timings

[Spray early in morning or evening or when overcast.](#)