

# **CABBAGE (TRANSPLANTED) - MEDIUM SOILS - LESS THAN 1.5% ORGANIC MATTER**

## General Information

### CABBAGE

(Transplanted only)

### Application Timing

- Apply pre-plant to stubble or soil surface in the fall or in the spring 60 days prior to planting or up to 72 hours after transplanting.
- If applying pre-emergence before transplant, make broadcast or banded applications.
- Early pre-plant applications can only be made in the states of CO, ID, MI, MN, MT, NE, ND, OR, SD, WA, WI, and WY.

### Important

- Read and follow all precautions, instructions, rotation crop guidelines, replanting instructions, and any other information listed on this label prior to use.
- Consult with university or extension weed management specialists for information on using Willowood Sulfentrazone 4SC with specific local varieties or cultivars of cabbage.

### Application Instructions

- Unless applying pre-plant incorporated, do not incorporate Willowood Sulfentrazone 4SC into the soil after application.  
Destroying the herbicide barrier by mechanically incorporating can allow weed escapes to occur.
- If applying pre-plant incorporated in the spring, mix thoroughly or shallowly incorporate Willowood Sulfentrazone 4SC into the soil prior to transplanting.
- If applying pre-emergence, applications before transplant can be broadcast or banded.
- Pre-emergence applications up to 72 hours after transplant should be a banded treatment in the row middles.
- Use rate is inversely dependent upon soil pH - use higher rates within the specified rate range with lower soil pH (<7.0) and lower rates within the specified rate range

with higher soil pH (>7.0).

#### Application Restrictions

- Do not incorporate into the soil deeper than 2".
- Do not apply Willowood Sulfentrazone 4SC to frozen or snow-covered soil.
- Do not apply more than 12.0 fl. oz. product per acre (0.375 lb. AI/A) per year (12-month period from when Willowood Sulfentrazone 4SC is first applied).
- Do not use on soils classified as "sand" (with <1% organic matter).

#### Limitations, Restrictions, and Exceptions

#### CABBAGE

(Transplanted only)

#### Application Timing

- Apply pre-plant to stubble or soil surface in the fall or in the spring 60 days prior to planting or up to 72 hours after transplanting.
- If applying pre-emergence before transplant, make broadcast or banded applications.
- Early pre-plant applications can only be made in the states of CO, ID, MI, MN, MT, NE, ND, OR, SD, WA, WI, and WY.

#### Important

- Read and follow all precautions, instructions, rotation crop guidelines, replanting instructions, and any other information listed on this label prior to use.
- Consult with university or extension weed management specialists for information on using Willowood Sulfentrazone 4SC with specific local varieties or cultivars of cabbage.

#### Application Instructions

- Unless applying pre-plant incorporated, do not incorporate Willowood Sulfentrazone 4SC into the soil after application. Destroying the herbicide barrier by mechanically incorporating can allow weed escapes to occur.
- If applying pre-plant incorporated in the spring, mix thoroughly or shallowly incorporate Willowood Sulfentrazone 4SC into the soil prior to transplanting.
- If applying pre-emergence, applications before transplant can be broadcast or banded.
- Pre-emergence applications up to 72 hours after transplant should be a banded

treatment in the row middles.

- Use rate is inversely dependent upon soil pH - use higher rates within the specified rate range with lower soil pH (<7.0) and lower rates within the specified rate range with higher soil pH (>7.0).

#### Application Restrictions

- Do not incorporate into the soil deeper than 2”.
- Do not apply Willowood Sulfentrazone 4SC to frozen or snow-covered soil.
- Do not apply more than 12.0 fl. oz. product per acre (0.375 lb. AI/A) per year (12-month period from when Willowood Sulfentrazone 4SC is first applied).
- Do not use on soils classified as “sand” (with <1% organic matter).

#### Method

[Broadcast/Foliar Air](#)

[Broadcast/Foliar Ground](#)

[Band application](#)

[Soil incorporation](#)

[Broadcast/Foliar Air](#)

[Broadcast/Foliar Ground](#)

[Band application](#)

[Soil incorporation](#)

[Broadcast/Foliar Air](#)

[Broadcast/Foliar Ground](#)

[Band application](#)

[Soil incorporation](#)

#### Rates

[field\\_rates 0](#)

- 

#### Restricted Entry Interval

12 hours

#### Soils

[Medium](#)

[Loam](#)

[Silt Loam](#)

[Silt](#)

[Sandy Clay Loam](#)

[Sandy Clay](#)

Timings

[Preemergence \(Crop\)](#)

[Preplant](#)

[Pretransplant](#)