

LETTUCE, LEAF CROPS

General Information

GENERAL INFORMATION

PHYTA-CAL QC can be tank mixed and applied with most commonly used insecticides and fungicides. DO NOT tank mix with growth hormone products, plant growth regulators, or foliar fertilizers unless specifically recommended.

DIRECTIONS FOR USE

Mix well before use. It is more effective to spray small amounts with greater frequency. DO NOT USE PHYTA-CAL QC WITH OTHER FOLIAR FERTILIZERS AS PRECIPITATION MAY OCCUR.

PHYTA-CAL QC is recommended on the following crops: alfalfa, almonds, apples, avocados, barley, all beans, broccoli, cabbage, carrots, cauliflower, celery, citrus, corn, grapes, lettuce, melons, milo, nectarines, peaches, peanuts, pears, pecans, peppers, potatoes, rice, sorghum, soybeans, strawberries, sugar beets, sugar cane, sweet corn, tomatoes, turnips, walnuts, watermelons, wheat, and most other crops.

Limitations, Restrictions, and Exceptions

Lettuce, Leaf Crops

Begin treatment after thinning and continue up to one week before harvest. Use 1 to 2 quarts per acre per application. Treat at 7 to 10 day intervals.

Method

[Spray](#)

Rates

[field rates 0](#)

-

Timings

[After thinning and continue up to one week before harvest.](#)