

# **COTTON, SOYBEANS, BEANS, TOMATOES, MELONS, CUCURBITS, PEANUTS, PEPPERS**

## General Information

### GENERAL INFORMATION

PHYTA-CAL QC can be tank mixed and applied with most commonly used insecticides and fungicides. DO NOT tank mix with growth hormone products, plant growth regulators, or foliar fertilizers unless specifically recommended.

### DIRECTIONS FOR USE

Mix well before use. It is more effective to spray small amounts with greater frequency. DO NOT USE PHYTA-CAL QC WITH OTHER FOLIAR FERTILIZERS AS PRECIPITATION MAY OCCUR.

PHYTA-CAL QC is recommended on the following crops: alfalfa, almonds, apples, avocados, barley, all beans, broccoli, cabbage, carrots, cauliflower, celery, citrus, corn, grapes, lettuce, melons, milo, nectarines, peaches, peanuts, pears, pecans, peppers, potatoes, rice, sorghum, soybeans, strawberries, sugar beets, sugar cane, sweet corn, tomatoes, turnips, walnuts, watermelons, wheat, and most other crops.

### Limitations, Restrictions, and Exceptions

Cotton, Soybeans, Beans, Tomatoes, Melons, Cucurbits, Peanuts, Peppers

Begin application when first flower appears. Spray enough water to achieve thorough coverage of the entire plant. Spray schedule may be discontinued when last fruit, boll, or pods are set. Use 2 to 4 quarts per treated acre at 7 to 10 day intervals.

### Method

[Spray](#)

### Rates

[field rates 0](#)

- 

### Timings

When first flower appears.