

WHEAT

General Information

Directions For Use

1. Stir well or agitate product prior to use.
2. Fill mix tank one-half to two-thirds full with water. Start and maintain constant agitation throughout entire mixing and treating process.
3. Add seed treatment materials, Color Coat Colorant and Flo Rite1197.
4. Bring the volume of slurry treatment mixture to the desired volume, with water.

Do not apply to grain used for food or feed purposes or processed for oil.

EXAMPLES:

When applied with neonicotinoid insecticide products at 0.25 mg ai /seed, apply Flo Rite 1197 at 0.46 mg /seed (2.0 fl oz/cwt based on 1400 kernels/pound).

When applied with neonicotinoid insecticide products at 0.5 mg ai /seed, apply Flo Rite 1197 at 0.92 mg/seed (4.0 fl oz/cwt based on 1400 kernels/pound).

When applied with neonicotinoid insecticide products at 1.25 mg ai/seed, apply Flo Rite 1197 at 2.2 mg /seed (9.5 fl oz/cwt based on 1400 kernels/pound).

-Suggested rates. Some variation may occur. Adjust the rates to obtain the desired coverage due to seed size, seed coat, conditioning equipment and total slurry.

Method

[Seed Treatment](#)

Rates

[field rates 0](#)

•

Timings

[N.A.](#)