

FOLIAR APPLICATION - DILUTE APPLICATIONS

General Information

General Information

BRANDT ORGANICS 5% Cu is recommended to prevent and correct copper deficiencies of field, row, vegetable, fruit, tree, and vine crops. Applied alone, with pesticides, or with other fertilizers, this product may be applied directly to the soil or to the foliage of growing crops via ground sprayers, aircraft or fertigation systems.

Mixing and Handling Instructions

Shake, stir, or swirl contents before using. Put 1/3 to 2/3 of total required water volume in tank. Add recommended amount of this product before adding pesticides, adjuvants, or other fertilizers. Agitate until thoroughly mixed, then fill tank with remainder of water. BRANDT ORGANICS 5% Cu is compatible with most pesticides and fertilizers. However, if the desired combination has not been previously used, check for chemical mixture compatibility using a jar test prior to application.

Limitations, Restrictions, and Exceptions

Rate Recommendations

FOLIAR APPLICATION: Actively growing plants with young leaves absorb foliarly applied BRANDT ORGANICS 5% Cu most efficiently. Immediate post-irrigation or post-rainfall application will enhance foliar uptake. Thorough coverage and wetting are needed for optimum foliar spray results.

Applications should be separated by a minimum of 7-14 days. AVOID foliar applications of this product when plants are under moisture stress. AVOID foliar applications to deciduous tree crops during bloom.

Dilute Applications: For rates higher than 2 quarts per acre, split applications of 2 quarts or less are preferred to minimize phytotoxicity.

Method

[Foliar application](#)

Rates

[field_rates 0](#)

-

Timings

[N.A.](#)