

SOIL APPLICATION - MILD DEFICIENCY

General Information

General Information

BRANDT ORGANICS 5% Cu is recommended to prevent and correct copper deficiencies of field, row, vegetable, fruit, tree, and vine crops. Applied alone, with pesticides, or with other fertilizers, this product may be applied directly to the soil or to the foliage of growing crops via ground sprayers, aircraft or fertigation systems.

Mixing and Handling Instructions

Shake, stir, or swirl contents before using. Put 1/3 to 2/3 of total required water volume in tank. Add recommended amount of this product before adding pesticides, adjuvants, or other fertilizers. Agitate until thoroughly mixed, then fill tank with remainder of water. BRANDT ORGANICS 5% Cu is compatible with most pesticides and fertilizers. However, if the desired combination has not been previously used, check for chemical mixture compatibility using a jar test prior to application.

Limitations, Restrictions, and Exceptions

SOIL APPLICATION: The lower rates are for field, row and vegetable crops. The higher rates are for fruit, tree, and vine crops.

Fertigation: Apply the recommended amount of product in split applications 1-2 weeks apart throughout the growing season.

Optimum rate of application will vary depending on treatment interval, soil properties (such as pH, organic matter content, texture), weather conditions, time of year, plant species and its nutrient requirements. For best results, follow soil/tissue test recommendation.

Method

[Soil application](#)

Rates

[field_rates 0](#)

-

Timings

[N.A.](#)