

# **POSTEMERGENT BROADLEAF WEED CONTROL IN CRANBERRY - OREGON**

## General Information

## Product Information

Stinger herbicide is a selective, postemergence herbicide for control of broadleaf weeds in apple, barley, oats and wheat not underseeded with a legume, canola (rapeseed), Christmas tree plantations, Conservation Reserve Program (CRP) acres, cottonwood/poplar and eucalyptus tree plantations, crambe, fallow cropland, field corn, garden beet, grasses grown for seed, Brassica, peppermint, popcorn, rangeland and permanent grass pastures, southern pine seedbeds in forest nurseries, spearmint, spinach, stone fruits, sugar beet, sweet corn, turnip, and non-cropland areas including fence rows, around farm buildings, and equipment pathways.

Stinger may be applied by aircraft on the following crops: canola (Rapeseed), crambe, spinach, and sugar beet.

Do not apply Stinger by aircraft to other labeled crops unless otherwise permitted by Dow Agro Sciences supplemental labeling.

Re-treatment is allowed, but do not apply more than the maximum allowable rate per crop growing season. An application to fallow cropland preceding or following an application to dryland small grains (wheat, barley or oats) is allowed, but is not allowed preceding or following an application to irrigated small grains.

## Restrictions

Use directions in Dow AgroSciences supplemental labeling may supersede directions or limitations in this labeling.

In California and New York, the maximum application rate for Stinger is 2/3 pint per acre per growing season. Do not exceed a cumulative amount of 2/3 pint of clopyralid [0.25 lb acid equivalent (a.e.)] per acre per crop year, unless specifically allowed.

Not for sale, use or distribution in Nassau and Suffolk Counties in New York State. Do not contaminate irrigation ditches or water used for irrigation or domestic purposes.

Do not use in greenhouses.

Chemigation: Do not apply this product through any type of irrigation system.

Do not spray pastures containing desirable forbs, especially legumes, unless injury can be tolerated.

Do not transfer livestock from treated grazing areas (or feeding of treated hay) to sensitive broadleaf crop areas without first allowing 7 days of grazing on an untreated pasture (or feeding of treated hay). If livestock are transferred within less than 7 days of grazing untreated pasture or eating untreated hay, urine and manure may contain enough clopyralid to cause injury to sensitive broadleaf plants.

Field Bioassay Instructions: In fields previously treated with this product, plant short test rows of the intended rotational crop across the original direction of application in a manner to sample field conditions, such as soil texture, soil pH, drainage, and any other variable that could affect the seed bed of the new crop. Field bioassay at any time prior to the planting of the intended rotational crop. Observe the test crop for herbicidal activity, such as poor stand (effect on seed germination) chlorosis (yellowing), necrosis (dead leaves or shoots), or stunting (reduced growth).

If herbicidal symptoms do not occur, the test crop can be grown. If there is apparent herbicidal activity, wait one year before repeating bioassay or plant only a labeled crop or crop listed in the table below for which the rotational interval has clearly been met.

#### Avoid Injury to Non-Target Plants

This product can affect susceptible broadleaf plants directly through foliage and indirectly by root uptake from treated soil. Therefore, do not apply Stinger directly to, or allow spray drift to come in contact with, vegetables, flowers, tomatoes, potatoes, beans, lentils, peas, alfalfa, sunflowers, soybeans, safflower, or other desirable broadleaf crops or ornamental plants or soil where sensitive crops will be planted the same season. (See Crop Rotation Intervals.)

Residues in Plants or Manure: Do not use plant residues, including hay or straw from treated areas, or manure or bedding straw from animals that have grazed or consumed forage from treated areas, for composting or mulching where susceptible plants may be grown the following season.

Do not spread manure from animals that have grazed or consumed forage or hay from treated areas on land used for growing susceptible broadleaf plants or apply such materials to land used for growing broadleaf crops, ornamentals, orchards, or other susceptible desirable plants. Plant materials or manure may contain enough clopyralid to cause injury to susceptible plant species. To promote herbicidal decomposition, plant residues should be evenly incorporated or burned. Breakdown

of clopyralid in crop residues or manure is more rapid under warm, moist soil conditions and may be enhanced by supplemental irrigation.

#### Avoid Movement of Treated Soil

Avoid conditions under which soil from treated areas may be moved or blown to areas containing susceptible plants. Wind-blown dust containing clopyralid may produce visible symptoms, such as epinasty (downward curving or twisting of leaf petioles or stems), when deposited on susceptible plants; however, serious injury is unlikely. To minimize potential movement of clopyralid on wind-blown dust, avoid treatment of powdery dry or light sandy soils until soil is settled by rainfall or irrigation or irrigate the treated soil shortly after application.

#### Application Directions

##### Application Timing

Apply to actively growing weeds. Extreme growing conditions, such as drought or near freezing temperatures prior to, at, or following application, may reduce weed control and increase the risk of crop injury at all stages of growth. Only weeds that have emerged at the time of application will be affected. If foliage is wet at the time of application, control may be decreased. Applications of Stinger are rainfast within 6 hours after application.

##### Application Rates

Generally, application rates at the lower end of the rate range will be satisfactory for young, succulent growth of susceptible weed species.

For less sensitive species, perennials, and under conditions where control is more difficult (plant stress conditions, such as, drought or extreme temperatures, dense weed stands and/or larger weeds), use a higher rate within the rate range. Weeds in fallow land or other areas where competition from crops is not present will generally require higher rates for control or suppression.

##### Spot Treatments

To prevent misapplication, apply spot treatments only with a calibrated boom or with hand sprayers according to directions provided below.

Hand Held Sprayers: Hand held sprayers may be used for spot applications. Care should be taken to apply the spray uniformly and at a rate equivalent to a broadcast application. Application rates in the table are based upon an area of 1000 sq ft. Mix the amount of Stinger (fl oz or mL) corresponding to the desired broadcast rate in 1 gallon or more of spray. To calculate the amount of Stinger required for

larger areas, multiply the table value (fl oz or mL) by the area to be treated in "thousands" of square feet, e.g., if the area to be treated is 3500 sq ft, multiply the table value by 3.5 (calc.  $3500 \div 1000 = 3.5$ ). An area of 1000 sq ft is approximately 10.5 x 10.5 yards (strides) in size.

#### Use of Adjuvants

Addition of surfactants, crop oils, or other adjuvants is not usually necessary when using Stinger. Adding a surfactant to the spray mixture may increase effectiveness on weeds but may reduce selectivity to the crop, particularly under conditions of plant stress. When an adjuvant is to be used with this product, Dow AgroSciences recommends the use of a Chemical Producers and Distributors Association certified adjuvant.

If an adjuvant is added to the spray solution, follow all manufacturer use guidelines.

#### Spray Coverage

Use sufficient spray volume to provide thorough coverage and a uniform spray pattern. Do not broadcast apply in less than 2 gallons total spray volume per acre. For best results, and to minimize spray drift, apply in a spray volume of 10 gallons or more per acre. As vegetative canopy and weed density increase, increase spray volume to obtain equivalent weed control. Use only nozzle types and spray equipment designed for herbicide application. To reduce spray drift, follow precautions under Avoid Injury to Non-Target Plants.

#### Limitations, Restrictions, and Exceptions

##### Use Restrictions

- Do not exceed a total of 2/3 pint per acre per year.
- Do not use surfactants when applying Stinger to cranberries.
- Do not spray once bud scales have separated and the growing point is visible.
- Do not apply within 5 feet of any water moving off or through the cranberry field.

Note: The total combined usage of Stinger from all types of applications must not exceed 2/3 pint per acre per year (0.25 lb ai/acre).

Broadcast Application Rates: Broadcast foliar application may be made at 1/4 to 2/3

pint per acre when cranberry plants are dormant or after terminal bud set. On new or nonbearing beds apply at 1/3 to 1/2 pint per acre. Use the lower rate in the rate range for young succulent growth for sensitive weed species. Use lower rates when application timing is close to bud break. Use the higher rate in the rate range for less sensitive weed species and under conditions where control is more difficult. Apply with a backpack sprayer or ground broadcast equipment in a total spray volume of 20 to 40 gallons of water per acre. The “timing window” for broadleaf weed control is based on the physiological state of the cranberry plant. This timing window begins when the cranberry vines enter dormancy in the fall and ends with budbreak in the spring (first emergence of bud expansion), when the crop becomes sensitive to application of Stinger. The time of budbreak is variety and weather dependent. An early or late spring can accelerate or delay budbreak, respectively. The ideal application window occurs when the weeds have emerged and have obtained sufficient canopy to allow treatment and when the cranberry plant is still dormant and tolerant to Stinger. Broadcast foliar applications that are made between budbreak and fruit set can cause plant injury.

**Hand-Held Sprayers:** Hand-held sprayers may be used for spot applications of Stinger if care is taken to apply the spray uniformly and at a rate equivalent to a broadcast application (example: 0.25 fl oz or 7.3 mL of Stinger per gallon of water is equal to the broadcast rate of 2/3 pint of Stinger per acre). To prevent misapplication, spot treatments should be applied with a calibrated boom or with hand sprayers according to the directions in the Hand-Held Sprayers section in the main label for Stinger for spot application instructions.

#### Specific Use Restrictions:

- Chemigation: Do not apply Stinger through any type of irrigation system.
- Preharvest Interval: Do not apply Stinger by any method within 60 days of harvest.
- Do not apply Stinger between budbreak and fruit set.
- Do not spray to runoff.
- Do not apply by aircraft
- Do not apply within 6 to 8 hours of expected rainfall or irrigation.
- Do not compost vegetation treated by broadcast application if compost will be used on sensitive plants.

#### Method

##### [Broadcast/Foliar Ground](#)

##### Pre-Harvest Interval

60 days

Rates

[field\\_rates 0](#)

[field\\_rates 1](#)

- 

Restricted Entry Interval

12 hours

Timings

[Postemergence \(Weed\)](#)

[When cranberry plants are dormant or after terminal bud set.](#)