

## **TREES WITH A 15 INCH OF TRUNK DIAMETER AT BREAST HEIGHT (D.B.H.) OR GREATER**

### General Information

#### NON-AGRICULTURAL USES

Do not apply Merit 2F Insecticide while bees are foraging. Do not apply Merit 2F Insecticide to plants that are flowering. Only apply after all flower petals have fallen off.

MERIT 2F Insecticide can be used as directed on sod farms, and on turfgrass, and ornamentals on such sites: residential home lawns, business and office complexes, shopping complexes, multi-family residential complexes, golf courses, airports, cemeteries, parks, playgrounds, and athletic fields.

MERIT 2F Insecticide can be used by soil application as directed on trees, shrubs, flowers, and groundcovers in sites: in and around the perimeter of industrial and commercial buildings and residential areas, and municipal, city, state, national, and private wooded and forested areas.

**RESTRICTION:** Not for use in commercial greenhouses, nurseries, or on grasses grown for seed, or on commercial fruit and nut trees.

**RESISTANCE:** Some insects are known to develop resistance to insecticides after repeated use. As with any insecticide, the use of this product should conform to resistance management strategies established for the use area.

MERIT 2F Insecticide contains a Group 4A insecticide. Insect biotypes with acquired or inherent resistance to Group 4A insecticides may eventually dominate the insect population if Group 4A insecticides are used repeatedly as the predominant method of control for targeted species.

### Limitations, Restrictions, and Exceptions

#### APPLICATION TO ORNAMENTALS

MERIT 2F Insecticide is for use on ornamentals in commercial and residential landscapes and interior plantscapes. MERIT 2F Insecticide is a systemic product and will be translocated upward into the plant system from root uptake. To assure

optimum effectiveness, the product must be placed where the growing portion of the target plant can absorb the active ingredient. The addition of a nitrogen containing fertilizer, where applicable, into the solution may enhance the uptake of the active ingredient. Application can be made by foliar application or soil applications; including soil injection, drenches, and broadcast sprays. Foliar applications offer locally systemic activity against insect pests.

When making soil applications to plants with woody stems, systemic activity will be delayed until the active ingredient is translocated throughout the plant. In some cases, this translocation delay could take 60 days or longer. For this reason, make application prior to anticipated pest infestation to achieve optimum levels of control.

**RESTRICTION:** Outdoor applications cannot exceed a total of 1.6 pints (0.4 lb of active ingredient) per acre per year.

#### ANT MANAGEMENT PROGRAMS

Use MERIT 2F Insecticide to control aphids, scale insects, mealybugs and other sucking pests on ornamentals to limit the honeydew available as a food source for ant populations. MERIT 2F Insecticide applications can be then be supplemented with residual sprays, bait placements or other ant control tactics to further reduce the pest population.

**Soil Injection: GRID SYSTEM:** Space holes on 2.5 foot centers, in a grid pattern, extending to the drip line of the tree. **CIRCLE SYSTEM:** Apply in holes evenly spaced in circles, (use more than one circle dependent upon the size of the tree) beneath the drip line of the tree extending in from that line. **BASAL SYSTEM:** Space injection holes evenly around the base of the tree trunk no more than 6 to 12 inches out from the base.

Mix required dosage in sufficient water to inject an equal amount of solution in each hole. Maintain a low pressure and use sufficient solution for distribution of the liquid into the treatment zone. For optimum control, keep the treated area moist for 7 to 10 days. Do not use less than 4 holes per tree or shrub.

Soil Drench: Uniformly apply the dosage in no less than 10 gallons of water per 1000 square feet as a drench around the base of the tree, directed to the root zone. Remove plastic or any other barrier that will stop solution from reaching the root zone.

For Control of Specified Borers: Application to trees already heavily infested may not prevent the eventual loss of the trees due to existing pest damage and tree stress.

**RESTRICTION:**

No Soil Injection Applications Allowed in Nassau or Suffolk Counties of New York.

Method

[Soil drench](#)

[Soil Injection](#)

Rates

[field\\_rates 0](#)

[field\\_rates 1](#)

- 

Restricted Entry Interval

12 hours

Exception: If the product is applied by drenching, the Worker Protection Standard, under certain circumstances, allows workers to enter the treated area if there will be no contact with anything that has been treated.

Timings

[N.A.](#)