

FOR USE ON CITRUS AND ORNAMENTALS TO CONTROL ASIAN CITRUS PSYLLID - CONTAINERIZED - TEXAS

General Information

NON-AGRICULTURAL USES

Do not apply Merit 2F Insecticide while bees are foraging. Do not apply Merit 2F Insecticide to plants that are flowering.

Only apply after all flower petals have fallen off.

MERIT 2F Insecticide can be used as directed on sod farms, and on turfgrass, and ornamentals on such sites: residential home lawns, business and office complexes, shopping complexes, multi-family residential complexes, golf courses, airports, cemeteries, parks, playgrounds, and athletic fields.

MERIT 2F Insecticide can be used by soil application as directed on trees, shrubs, flowers, and groundcovers in sites: in and around the perimeter of industrial and commercial buildings and residential areas, and municipal, city, state, national, and private wooded and forested areas.

RESTRICTION: Not for use in commercial greenhouses, nurseries, or on grasses grown for seed, or on commercial fruit and nut trees.

RESISTANCE: Some insects are known to develop resistance to insecticides after repeated use. As with any insecticide, the use of this product should conform to resistance management strategies established for the use area.

MERIT 2F Insecticide contains a Group 4A insecticide. Insect biotypes with acquired or inherent resistance to Group 4A insecticides may eventually dominate the insect population if Group 4A insecticides are used repeatedly as the predominant method of control for targeted species.

Limitations, Restrictions, and Exceptions

Rate: 0.025 fl oz (0.75 ml) per ft³ of container media

Restrictions:

Preharvest interval (PHI): 0 days

Recommended applications:

Determine volume of container and calculate dosage necessary to treat container. Apply calculated dosage of MERIT 2F Insecticide per container as a soil drench. Use sufficient volume to ensure thorough and uniform distribution throughout the media without loss of gravitational water from the container. Retreat if necessary.

Keep children and pets off treated area until dry.

Method

[Soil drench](#)

Rates

[field_rates 0](#)

[field_rates 1](#)

-

Restricted Entry Interval

12 hours

Exception: If the product is applied by drenching, the Worker Protection Standard, under certain circumstances, allows workers to enter the treated area if there will be no contact with anything that has been treated.

Timings

[N.A.](#)