

ORNAMENTALS - CONTAINER GROWN ORNAMENTALS

Limitations, Restrictions, and Exceptions

DIRECTIONS FOR APPLICATION TO ORNAMENTALS

Drench, Crown and Surface Spray Application: DISARM 480 SC may be applied to control soilborne, seedling, and crown diseases of production ornamentals (greenhouse, shadehouse, container grown and field grown) as a preventative, drench, crown or surface spray treatment prior to infection. Good coverage of the pre-infection area (root zone, root ball, crown, etc.) is necessary for satisfactory control. Make applications prior to infection as healthy roots are necessary to optimize product uptake systemic translocation and disease protection.

DISARM 480 SC may be applied as drench to container grown ornamentals using 0.15–0.6 fl oz/100 gallons of water. Thoroughly wet the root zone of the plants with the solution using up to 1–2 pt per sq ft of surface area. If a drench application is not feasible, use 0.4–1.6 fl oz/1,000 sq ft of growing area in sufficient water to provide uniform coverage and follow with enough irrigation to completely wet the root zone of the plants. Use of the higher rate drench is limited to one application per year.

Restrictions: Do not apply more than 2.13 lbs ai per acre per year.

Method

[Drench](#)

Rates

[field rates 0](#)

•

Timings

[Prior to infection as healthy roots are necessary to optimize product uptake systemic translocation and disease protection.](#)