

CHRYSANTHEMUMS (CUTTINGS) - PRE-PLANT DIP

General Information

GENERAL INFORMATION

B-NINE WSG is a plant growth regulator for use on container grown ornamental plants. When applied to the plant foliage B-NINE WSG passes into the leaf, then moves within the plant to the growing points to reduce internode elongation. Treated plants will develop a shorter, more compact growth habit when compared to untreated plants.

FOR USE ONLY ON ORNAMENTALS GROWN IN COMMERCIAL OR RESEARCH GREENHOUSES, SHADEHOUSES AND NURSERIES. IN PRODUCTION AREAS NOT UNDER COVER, USE IS RESTRICTED TO CONTAINERIZED ORNAMENTALS¹

APPLICATION NOTES

The recommended application volume is 1 gallon of B-NINE WSG spray solution to cover 200 square feet of area. This will provide foliage coverage just to the point of runoff.

Plants should be well watered prior to application of B-NINE WSG. Wilted plants do not readily absorb B-NINE WSG and poor results might be obtained if application is made to wilted plants.

The plant foliage should be dry when sprays are applied. Do not syringe or overspray the foliage for 18 to 24 hours after applying B-NINE WSG to allow the chemical to enter the plant.

Two applications of B-NINE WSG at 1/2 to 2/3 the label rates may give better control than a single application of the total recommended rate.

¹Use on bed-grown or containerized ornamentals without impervious floors not permitted by the Arizona Department of Environmental Quality.

USE AND RATE RECOMMENDATIONS BY CROP

Although the use of B-NINE WSG has been found to be acceptable for the specific plants listed in this label, be sure to read and fully understand the section on

DETERMINING PLANT SAFETY before applying to large numbers of plants.

Limitations, Restrictions, and Exceptions

CHRYSANTHEMUMS (CUTTINGS)

At certain times of the year or with certain tall or long internoded varieties, the conventional spray applications described above may not be adequate. Additional applications of B-NINE WSG may be useful as pre-plant dips and post-plant sprays as follows.

Pre-plant dip: - Use 0.10% (1000 ppm):

ROOTED CUTTING: Hold a handful of cuttings by the root system; dip in the B-NINE WSG solution just long enough to thoroughly wet all leaves and stems, remove and pot. Allow the foliage to dry before watering-in. Normally, by the time you are at the end of a bed, you can start watering-in those plants first potted.

UNROOTED CUTTING: Remove unrooted cuttings from the box. Dip stem in the B-NINE WSG solution, remove and place in flat. Cover with cheesecloth to avoid dehydration and place in the cooler overnight. Stick the next day and go on to a normal misting program.

Method

[Dip](#)

Rates

[field rates 0](#)

[field rates 1](#)

•

Restricted Entry Interval

24 hours

Timings

[Pre-plant.](#)